

Existence of Continuous and Càdlàg Versions for Cylindrical Processes in the Dual of a Nuclear Space

C. A. Fonseca-Mora

School of Mathematics and Statistics, The University of Sheffield, Hicks Building,
Hounsfield Road, S3 7RH, United Kingdom,

and

Escuela de Matemática, Universidad de Costa Rica, San Pedro de Montes de Oca,
San José, Apartado Postal 11501-2060, Costa Rica.

Email: christianandres.fonseca@ucr.ac.cr

Abstract

Let Φ be a nuclear space and let Φ'_β denote its strong dual. In this paper we introduce sufficient conditions for a cylindrical process in Φ' to have a version that is a Φ'_β -valued continuous or càdlàg process. We also establish sufficient conditions for the existence of such a version taking values and having finite moments in a Hilbert space continuously embedded in Φ'_β . Finally, we apply our results to the study of properties of cylindrical martingales in Φ' .

2010 Mathematics Subject Classification: 60B11, 60G20, 60G17, 28C20.

Key words and phrases: Cylindrical stochastic processes, dual of a nuclear space, continuous and càdlàg versions, cylindrical martingales.

1 Introduction

Probability theory in infinite dimensional spaces has been an area under intensive development since the 1960s. There are several monographs devoted to the study of stochastic processes and measures defined on these spaces; we can cite for example [2, 5, 12, 13, 19, 20, 34, 36].

Among the range of topics within the theory of probability on infinite dimensional spaces is the study of cylindrical probability measures, that is, finitely additive set functions that have “projections” on finite dimensional spaces that are probability measures. In general, cylindrical probability measures do not necessarily extend to a countably additive probability measures; the investigation of this problem led to the celebrated discoveries of Minlos (see [22]) and Sazonov (see [32]) in the context of duals of countably Hilbertian nuclear spaces and of Hilbert spaces respectively, which establish necessary and sufficient conditions in terms of the continuity of the Fourier transform of a cylindrical measure for this extension to be possible.

Associated to the concept of cylindrical measures is that of cylindrical random variables. These are similar to the usual random variables but their laws are cylindrical, rather than bona fide, probability measures. Recently, cylindrical processes, i.e. collections of cylindrical random variables indexed by time, are increasingly attracting attention because they appear naturally as the driving noise of stochastic partial differential equations (see e.g. [1, 17, 27, 29, 30, 31, 37]).

Parallel to the problem of finding conditions for the extension of cylindrical probability measures to countably additive probability measures is the problem of

finding a random variable Y in the underlying space that is a version of a given cylindrical random variable X , where here “version” means that Y coincides almost surely with X when evaluated with respect to any element of the dual space. In the case of the dual of a nuclear space, the regularization theorem of Itô and Nawata (see [14]) established necessary and sufficient conditions for the existence of such a version for a cylindrical random variable. As a consequence of the regularization theorem we can establish sufficient conditions for the existence of an stochastic process that is a version of a given cylindrical process.

In [24], Mitoma addressed the more specialized problem of establishing sufficient conditions for an stochastic process taking values in the dual of a nuclear Fréchet space to have a continuous or a right-continuous with left limits version. There are several applications for this result. Some examples are the study of self-intersection local times of density processes in $\mathcal{S}'(\mathbb{R}^d)$ (see [4]) and the study of regularity properties of the limit process of many-server queueing systems (see [18]). By using different techniques, several authors extended Mitoma’s work by establishing different conditions on the nuclear space and on the stochastic processes (see [8, 10, 21]).

In this paper we carried out a further extension by showing that under some natural conditions a cylindrical processes in the dual of a nuclear space (no other conditions are assumed on the space) has a continuous or a right-continuous with left limits version. To the extend of our knowledge our result is the more general result of its type that can be found on the literature. Moreover, our result is a natural generalization of the regularization theorem of Itô and Nawata for cylindrical random variables.

As well as from proving this result, we also introduce some new results on the existence of continuous and càdlàg versions taking values in a Hilbert space continuously embedded in the dual of a nuclear space. Then we apply our results to the existence of continuous and càdlàg versions for stochastic processes and to the study of martingales taking values in the dual of a nuclear space.

We remark that we have successfully applied the results obtained in this paper to other problems, for example to prove the Lévy-Itô decomposition for Lévy processes taking values in the strong dual of a reflexive nuclear space and to the introduction of a new theory of stochastic integration for operator-valued processes with respect to Lévy processes on the dual of a nuclear space; both of these are furthermore applied to the study of stochastic evolution equations driven by Lévy noise in duals of nuclear spaces (see [9]). These results will be published elsewhere.

The organization of the paper is as follows. In Section 2 we review the basic concepts on nuclear spaces and of random processes defined on its strong dual that we will need throughout this paper. Section 3 is devoted to the proof of our main result (Theorem 3.2). In Section 4 we prove several modifications of our result for the existence of continuous and càdlàg versions taking values in a Hilbert space continuously embedded in the dual of a nuclear space. Finally, in Section 5 we apply our results to study cylindrical martingales in the dual of a nuclear space.

2 Preliminaries

2.1 Nuclear Spaces

In this section we introduce our notation and review some of the key concepts that we will need throughout this paper. For detailed information on locally convex spaces the reader is referred to [15, 33, 35].

Let Φ be a locally convex space (over $\mathbb{R}$ or $\mathbb{C}$). If p is a continuous semi-norm on Φ and $r > 0$, the closed ball of radius r of p given by $B_p(r) = \{\phi \in \Phi : p(\phi) \leq r\}$ is a closed, convex, balanced neighborhood of zero in Φ .

A continuous semi-norm (respectively a norm) p on Φ is called *Hilbertian* if $p(\phi)^2 = Q(\phi, \phi)$, for all $\phi \in \Phi$, where Q is a symmetric, non-negative bilinear form (respectively inner product) on $\Phi \times \Phi$. Let Φ_p be the Hilbert space that corresponds to the completion of the pre-Hilbert space $(\Phi/\ker(p), \tilde{p})$, where $\tilde{p}(\phi + \ker(p)) = p(\phi)$ for each $\phi \in \Phi$. The quotient map $\Phi \rightarrow \Phi/\ker(p)$ has a unique continuous linear extension $i_p : \Phi \rightarrow \Phi_p$.

Let q be another continuous Hilbertian semi-norm on Φ for which $p \leq q$. In this case, $\ker(q) \subseteq \ker(p)$. Moreover, the inclusion map from $\Phi/\ker(q)$ into $\Phi/\ker(p)$ is linear and continuous, and therefore it has a unique continuous extension $i_{p,q} : \Phi_q \rightarrow \Phi_p$. Furthermore, we have the following relation: $i_p = i_{p,q} \circ i_q$.

We denote by Φ' the topological dual of Φ and by $f[\phi]$ the canonical pairing of elements $f \in \Phi'$, $\phi \in \Phi$. We denote by Φ'_β the dual space Φ' equipped with its *strong topology* β , i.e. β is the topology on Φ' generated by the family of semi-norms $\{\eta_B\}$, where for each $B \subseteq \Phi'$ bounded we have $\eta_B(f) = \sup\{|f[\phi]| : \phi \in B\}$ for all $f \in \Phi'$. If p is a continuous Hilbertian semi-norm on Φ , then we denote by Φ'_p the Hilbert space dual to Φ_p . The dual norm p' on Φ'_p is given by $p'(f) = \sup\{|f[\phi]| : \phi \in E, p(\phi) \leq 1\}$ for all $f \in \Phi'_p$. Moreover, the dual operator i'_p corresponds to the canonical inclusion from Φ'_p into Φ'_β and it is linear and continuous.

Let p and q be continuous Hilbertian semi-norms on Φ such that $p \leq q$. The space of continuous linear operators (respectively Hilbert-Schmidt operators) from Φ_q into Φ_p is denoted by $\mathcal{L}(\Phi_q, \Phi_p)$ (respectively $\mathcal{L}_2(\Phi_q, \Phi_p)$) and the operator norm (respectively Hilbert-Schmidt norm) is denoted by $\|\cdot\|_{\mathcal{L}(\Phi_q, \Phi_p)}$ (respectively $\|\cdot\|_{\mathcal{L}_2(\Phi_q, \Phi_p)}$). We employ an analogue notation for operators between the dual spaces Φ'_p and Φ'_q .

Among the many equivalent definitions of a nuclear space (see [28, 35]), the following is the most useful for our purposes.

Definition 2.1. A (Hausdorff) locally convex space $(\Phi, \mathcal{T})$ is called *nuclear* if its topology $\mathcal{T}$ is generated by a family Π of Hilbertian semi-norms such that for each $p \in \Pi$ there exists $q \in \Pi$, satisfying $p \leq q$ and the canonical inclusion $i_{p,q} : \Phi_q \rightarrow \Phi_p$ is Hilbert-Schmidt.

Example 2.2. The following locally convex spaces are nuclear (see [35], Chapter 51 and [28], Chapter 6): the (Schwartz) space of rapidly decreasing functions $\mathcal{S}(\mathbb{R}^d)$ and the space of tempered distributions $\mathcal{S}'(\mathbb{R}^d)$; the space $\mathcal{D}(X)$ of test functions on X and the space of distributions $\mathcal{D}'(X)$ (X : open subset of $\mathbb{R}^d$); the space $\mathcal{C}^\infty(U)$ of infinitely differentiable real or complex functions on U (U : open subset of $\mathbb{R}^d$) and the space $\mathcal{H}(G)$ of holomorphic functions on G (G : open subset of $\mathbb{C}^n$).

Definition 2.3. Let Φ be a vector space and let Π be a family of Hilbertian semi-norms on Φ such that for each $p \in \Pi$ the Hilbert space Φ_p is separable. The locally convex topology $\mathcal{T}$ on Φ generated by the family Π is called *multi-Hilbertian* (respectively *countably Hilbertian* if Π is countable). The space $(\Phi, \mathcal{T})$ is called a *multi-Hilbertian space* (respectively *countably Hilbertian*).

If $(\Phi, \mathcal{T})$ is a nuclear space and p is a continuous Hilbertian semi-norm on Φ , the Hilbert space Φ_p is separable (see [28], Proposition 4.4.9 and Theorem 4.4.10, p.82). Therefore, every nuclear space is multi-Hilbertian.

Let $(\Phi, \mathcal{T})$ be a multi-Hilbertian space and let $\{p_n\}_{n \in \mathbb{N}}$ be an increasing sequence of continuous Hilbertian semi-norms on $(\Phi, \mathcal{T})$. Denote by θ the countably Hilbertian topology on Φ generated by the semi-norms $\{p_n\}_{n \in \mathbb{N}}$. The topology θ is weaker than $\mathcal{T}$. We denote by Φ_θ the space (Φ, θ) . Some properties of the space Φ_θ are listed below:

Proposition 2.4. *In the above notation,*

- (1) Φ_θ is a separable pseudo-metrizable locally convex space and $d : \Phi \rightarrow \mathbb{R}$ given by $d(\phi) = \sum_{n \in \mathbb{N}} 2^{-n} [p_n(\phi)/(1 + p_n(\phi))]$ for $\phi \in \Phi$ is a pseudo-metric for Φ_θ .

(2) The completion $\widetilde{\Phi}_\theta$ of Φ_θ is separable, complete, pseudo-metrizable space and it is furthermore a Baire space.

(3)

$$(\widetilde{\Phi}_\theta)' = \Phi'_\theta = \bigcup_{n \in \mathbb{N}} \Phi'_{p_n}. \quad (2.1)$$

Proof. (1) The fact that Φ_θ is pseudo-metrizable and that d is a pseudo-metric is a consequence that its topology is generated by a countable number of semi-norms (see [26], Theorem 5.6.1, p.123). The fact that Φ_θ is separable is a consequence of the fact that $\{p_n\}_{n \in \mathbb{N}}$ generates the topology θ and that Φ_{p_n} is separable for each $n \in \mathbb{N}$ (see [28], Theorem 4.4.10, p.82-3).

(2) The completion $\widetilde{\Phi}_\theta$ of Φ_θ is again a pseudo-metrizable locally convex space (see [26], Theorem 3.7.1, p.60) and hence by (an extension of) the Baire category theorem (see [26], Theorem 11.7.2, p.393) it follows that $\widetilde{\Phi}_\theta$ is a Baire space.

(3) It is well-know that the dual space of Φ_θ is equal to $\bigcup_{n \in \mathbb{N}} \Phi'_{p_n}$. On the other hand, the dual space $(\widetilde{\Phi}_\theta)'$ of $\widetilde{\Phi}_\theta$ can be identified (algebraically) with the dual space Φ'_θ of Φ_θ (see [15], Corollary 4, Section 3.4, p.63), thus we obtain (2.1). $\square$

Remark 2.5. Even if $(\Phi, \mathcal{T})$ is Hausdorff this does not in general implies that Φ_θ is Hausdorff. However, if Φ_θ is Hausdorff the pseudo-metric d defined in Proposition 2.4 is a metric and hence Φ_θ is metrizable (see [15], Theorem 1, Section 2.8, p.40). Moreover, in that case $\widetilde{\Phi}_\theta$ is a Fréchet space and it is (isomorphic to) the projective limit $\text{proj}_{n \in \mathbb{N}} \Phi_{p_n}$ (see [15], Corollary 7, Section 3.4, p.63).

2.2 Cylindrical and Stochastic Processes

Assumption 2.6. Unless otherwise specified, in this section Φ will always denote a multi-Hilbertian space over $\mathbb{R}$.

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a complete probability space. We denote by $L^0(\Omega, \mathcal{F}, \mathbb{P})$ the space of equivalence classes of real-valued random variables defined on $(\Omega, \mathcal{F}, \mathbb{P})$. We always consider the space $L^0(\Omega, \mathcal{F}, \mathbb{P})$ equipped with the topology of convergence in probability and in this case it is a complete, metrizable, topological vector space, but is not in general locally convex (see e.g. [2]).

A Borel measure μ on Φ'_β is called a *Radon measure* if for every $\Gamma \in \mathcal{B}(\Phi'_\beta)$ and $\epsilon > 0$, there exist a compact set $K_\epsilon \subseteq \Gamma$ such that $\mu(\Gamma \setminus K_\epsilon) < \epsilon$. In general not every Borel measure on Φ is Radon, however, when Φ is a Fréchet nuclear space or a countable inductive limit of Fréchet nuclear spaces, then every Borel measure on Φ'_β is a Radon measure (see Corollary 1.3 of Dalecky and Fomin [5], p.11).

For any $n \in \mathbb{N}$ and any $\phi_1, \dots, \phi_n \in \Phi$, we define a linear map $\pi_{\phi_1, \dots, \phi_n} : \Phi' \rightarrow \mathbb{R}^n$ by

$$\pi_{\phi_1, \dots, \phi_n}(f) = (f[\phi_1], \dots, f[\phi_n]), \quad \forall f \in \Phi'. \quad (2.2)$$

The map $\pi_{\phi_1, \dots, \phi_n}$ is clearly linear and continuous. Let M be a subset of Φ . A subset of Φ' of the form

$$\mathcal{Z}(\phi_1, \dots, \phi_n; A) = \{f \in \Phi' : (f[\phi_1], \dots, f[\phi_n]) \in A\} = \pi_{\phi_1, \dots, \phi_n}^{-1}(A) \quad (2.3)$$

where $n \in \mathbb{N}$, $\phi_1, \dots, \phi_n \in M$ and $A \in \mathcal{B}(\mathbb{R}^n)$ is called a *cylindrical set* based on M . The set of all the cylindrical sets based on M is denoted by $\mathcal{Z}(\Phi', M)$. It is an algebra but if M is a finite set then it is a σ -algebra. The σ -algebra generated by $\mathcal{Z}(\Phi', M)$ is denoted by $\mathcal{C}(\Phi', M)$ and it is called the *cylindrical σ -algebra* with respect to (Φ', M) . If $M = \Phi$, we write $\mathcal{Z}(\Phi') = \mathcal{Z}(\Phi', \Phi)$ and $\mathcal{C}(\Phi') = \mathcal{C}(\Phi', \Phi)$. One can easily see from (2.3) that $\mathcal{Z}(\Phi'_\beta) \subseteq \mathcal{B}(\Phi'_\beta)$. Therefore, $\mathcal{C}(\Phi'_\beta) \subseteq \mathcal{B}(\Phi'_\beta)$. In general this inclusion is strict but if Φ is separable (for example if it is a countably

Hilbertian space) then $\mathcal{C}(\Phi'_\beta) = \mathcal{B}(\Phi'_\beta)$ (see Lemma 4.1 in Mitoma, Okada and Okazaki [25]).

A function $\mu : \mathcal{Z}(\Phi') \rightarrow [0, \infty]$ is called a *cylindrical measure* on Φ' , if for each finite subset $M \subseteq \Phi'$ the restriction of μ to $\mathcal{C}(\Phi', M)$ is a measure. A cylindrical measure μ is called *finite* if $\mu(\Phi') < \infty$ and a *cylindrical probability measure* if $\mu(\Phi') = 1$. The complex-valued function $\hat{\mu} : \Phi \rightarrow \mathbb{C}$ defined by

$$\hat{\mu}(\phi) = \int_{\Phi'} e^{if[\phi]} \mu(df) = \int_{-\infty}^{\infty} e^{iz} \mu_\phi(dz), \quad \forall \phi \in \Phi,$$

where for each $\phi \in \Phi$, $\mu_\phi := \mu \circ \pi_\phi^{-1}$, is called the *characteristic function* of μ . In general, a cylindrical measure on Φ' does not extend to a Borel measure on Φ'_β . However, necessary and sufficient conditions for this can be given in terms of the continuity of its characteristic function by means of the Minlos theorem (see [5], Theorem 1.3, Chapter III, p.88).

Theorem 2.7 (Minlos theorem). *Let Φ be a nuclear space. For a function $F : \Phi \rightarrow \mathbb{C}$ to be the characteristic function of a Radon probability measure on Φ'_β it is sufficient, and necessary if Φ is barrelled, that it be positive definite, continuous at zero and satisfies $F(0) = 1$.*

A *cylindrical random variable* in Φ' is a linear map $X : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$. If $Z = \mathcal{Z}(\phi_1, \dots, \phi_n; A)$ is a cylindrical set, for $\phi_1, \dots, \phi_n \in \Phi$ and $A \in \mathcal{B}(\mathbb{R}^n)$, let

$$\mu_X(Z) := \mathbb{P}((X[\phi_1], \dots, X[\phi_n]) \in A) = \mathbb{P} \circ X^{-1} \circ \pi_{\phi_1, \dots, \phi_n}^{-1}(A).$$

The map μ_X is called the *cylindrical distribution* of X and it is cylindrical probability measure on Φ' .

If X is a cylindrical random variable in Φ' , the *characteristic function* of X is defined to be the characteristic function $\hat{\mu}_X : \Phi \rightarrow \mathbb{C}$ of its cylindrical distribution μ_X . Therefore, $\hat{\mu}_X(\phi) = \mathbb{E}e^{iX(\phi)}$, $\forall \phi \in \Phi$. Also, we say that X is *n-integrable* if $\mathbb{E}(|X(\phi)|^n) < \infty$, $\forall \phi \in \Phi$.

Let X be a Φ'_β -valued random variable, i.e. $X : \Omega \rightarrow \Phi'_\beta$ is a $\mathcal{F}/\mathcal{B}(\Phi'_\beta)$ -measurable map. We denote by μ_X the distribution of X , i.e. $\mu_X(\Gamma) = \mathbb{P}(X \in \Gamma)$, $\forall \Gamma \in \mathcal{B}(\Phi'_\beta)$, and it is a Borel probability measure on Φ'_β . For each $\phi \in \Phi$ we denote by $X[\phi]$ the real-valued random variable defined by $X[\phi](\omega) := X(\omega)[\phi]$, for all $\omega \in \Omega$. Then, the mapping $\phi \mapsto X[\phi]$ defines a cylindrical random variable. Therefore, the above concepts of characteristic function and integrability can be analogously defined for Φ'_β -valued random variables in terms of the cylindrical random variable they determines.

Very important for our forthcoming arguments is the following class of Φ'_β -valued random variables introduced by Itô and Nawata in [14].

Definition 2.8. A Φ'_β -valued random variable X is called *regular* if there exists a weaker countably Hilbertian topology θ on Φ such that $\mathbb{P}(\omega : X(\omega) \in \Phi'_\theta) = 1$.

If X is a cylindrical random variable in Φ' , a Φ'_β -valued random variable Y is called a *version* of X if for every $\phi \in \Phi$, $X(\phi) = Y[\phi]$ $\mathbb{P}$ -a.e. A sufficient condition for the existence of a regular version is given in the following result due to Itô and Nawata (see [14]):

Theorem 2.9 (Regularization theorem). *Let X be a cylindrical random variable in Φ' such that $X : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is continuous. Then, X has a unique (up to equivalence) Φ'_β -valued regular version.*

The following results establish alternative characterizations for regular random variables.

Theorem 2.10. *Let X be a Φ'_β -valued random variable. Consider the statements:*

(1) X is regular.

(2) The map $X : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$, $\phi \mapsto X[\phi]$ is continuous.

(3) The distribution μ_X of X is a Radon probability measure.

One has the implications: (1) $\Rightarrow$ (2) and if Φ is nuclear, then (2) $\Rightarrow$ (1) and (2) $\Rightarrow$ (3). Moreover, if Φ is barrelled, then (3) $\Rightarrow$ (1).

Proof. (1) $\Rightarrow$ (2): Let θ be a countably Hilbertian topology on Φ as in Definition 2.8. As the topology θ is weaker than the nuclear topology on Φ , it is sufficient to show that the map $\phi \mapsto X[\phi]$ is continuous from Φ_θ into $L^0(\Omega, \mathcal{F}, \mathbb{P})$. To show this, note that if $\{\phi_n\}_{n \in \mathbb{N}}$ is a sequence converging to ϕ in Φ_θ , then the fact that $\mathbb{P}(\omega : X(\omega) \in \Phi'_\theta) = 1$ shows that $X[\phi_n] \rightarrow X[\phi]$ $\mathbb{P}$ -a.e. Therefore $|X[\phi_n] - X[\phi]| \rightarrow 0$ in probability. Hence, because Φ_θ is first-countable (this is because it is pseudo-metrizable, see Proposition 2.4) it then follows that the map $\phi \mapsto X[\phi]$ is continuous from Φ_θ into $L^0(\Omega, \mathcal{F}, \mathbb{P})$.

(2) $\Rightarrow$ (1): As the map $\phi \mapsto X[\phi]$ is continuous, because Φ is nuclear the regularization theorem (Theorem 2.9) shows that X has a regular version. But this clearly implies that X is also regular.

(2) $\Rightarrow$ (3): The assumption that $\phi \mapsto X[\phi]$ is continuous implies that the characteristic function of X is continuous. If Φ is nuclear the Minlos theorem (Theorem 2.7) shows that the distribution μ_X of X is a Radon probability measure.

(3) $\Rightarrow$ (1): Assume that Φ is barrelled and that μ_X is a Radon measure. Let $\{\epsilon_n\}_{n \in \mathbb{N}}$ be a decreasing sequence of positive real numbers converging to zero. For every $n \in \mathbb{N}$, because μ_X is a Radon probability measure there exists a compact subset K_n of Φ'_β such that $\mu_X(K_n) > 1 - \epsilon_n$.

Because Φ is barrelled and each K_n is bounded, there exists an increasing sequence of continuous Hilbertian semi-norms $\{p_n\}_{n \in \mathbb{N}}$ on Φ such that for every $n \in \mathbb{N}$, $K_n \subseteq B_{p_n}(1)^0$, where $B_{p_n}(1)^0$ is the polar set of $B_{p_n}(1)$ (see [33], Result 5.2, Chapter IV, p.141). Then, as $B_{p_n}(1)^0$ is the unit ball of the Hilbert space Φ'_{p_n} , it follows that

$$\mathbb{P}(X \in \Phi'_{p_n}) = \mu_X(\Phi'_{p_n}) \geq \mu_X(B_{p_n}(1)^0) \geq \mu_X(K_n) > 1 - \epsilon_n.$$

Hence, as $\epsilon_n \rightarrow 0$ it follows from the above inequality that $\mathbb{P}(X \in \bigcup_{n \in \mathbb{N}} \Phi'_{p_n}) = 1$. Then, if θ is the countably Hilbertian topology generated by the semi-norms $\{p_n\}_{n \in \mathbb{N}}$ it follows from (2.1) that $\mathbb{P}(\omega : X(\omega) \in \Phi'_\theta) = 1$ and hence X is a regular random variable. $\square$

The following is an useful property of regular random variables.

Proposition 2.11. *Let X, Y be Φ'_β -valued random regular variables. Then, $X = Y$ $\mathbb{P}$ -a.e. if and only if for all $\phi \in \Phi$, $X[\phi] = Y[\phi]$ $\mathbb{P}$ -a.e.*

Proof. We only have to prove the sufficiency. Assume X and Y are regular and let $\{p_n\}_{n \in \mathbb{N}}$ be a sequence of Hilbertian semi-norms on Φ such that Definition 2.8 is satisfied for both X and Y . If θ is the countably Hilbertian topology on Φ determined by the semi-norms $\{p_n\}_{n \in \mathbb{N}}$, then $\mathbb{P}(\Omega_\theta) = 1$, where $\Omega_\theta = \{\omega \in \Omega : X(\omega) \in \Phi'_\theta, Y(\omega) \in \Phi'_\theta\}$.

Because Φ_θ is separable, there exists a countable subset $\{\phi_j : j \in \mathbb{N}\}$ of Φ that is dense in Φ_θ . For every $j \in \mathbb{N}$, it follows from our hypothesis that $\mathbb{P}(\Omega_j) = 1$, where $\Omega_j = \{\omega \in \Omega : X(\omega)[\phi_j] = Y(\omega)[\phi_j]\}$. Let $\Gamma = \Omega_\theta \cap \bigcap_{j \in \mathbb{N}} \Omega_j$. Then, we have $\mathbb{P}(\Gamma) = 1$.

Fix $\omega \in \Gamma$ and let $\phi \in \Phi$. Then, there exists a sequence $\{\phi_{j_k}\}_{k \in \mathbb{N}} \subseteq \{\phi_j : j \in \mathbb{N}\}$ that converges to ϕ in Φ_θ . Therefore, as $X(\omega), Y(\omega) \in \Phi'_\theta$ and because $X(\omega)[\phi_{j_k}] = Y(\omega)[\phi_{j_k}]$ for all $k \in \mathbb{N}$, it follows that

$$X(\omega)[\phi] = \lim_{k \rightarrow \infty} X(\omega)[\phi_{j_k}] = \lim_{k \rightarrow \infty} Y(\omega)[\phi_{j_k}] = Y(\omega)[\phi].$$

As the above is true for any $\phi \in \Phi$, it follows that $X(\omega) = Y(\omega)$ for every $\omega \in \Gamma$. Therefore, $X = Y$ $\mathbb{P}$ -a.e. $\square$

Let $J = [0, \infty)$ or $J = [0, T]$ for some $T > 0$. We say that $X = \{X_t\}_{t \in J}$ is a *cylindrical process* in Φ' if X_t is a cylindrical random variable, for each $t \in J$. Clearly, any Φ'_β -valued stochastic processes $X = \{X_t\}_{t \in J}$ defines a cylindrical process under the prescription: $X[\phi] = \{X_t[\phi]\}_{t \in J}$, for each $\phi \in \Phi$. We will say that it is the *cylindrical process determined by X* .

A Φ'_β -valued processes $Y = \{Y_t\}_{t \in J}$ is said to be a Φ'_β -valued *version* of the cylindrical process $X = \{X_t\}_{t \in J}$ on Φ' if for each $t \in J$, Y_t is a Φ'_β -valued version of X_t .

A Φ'_β -valued stochastic process X is *continuous* (respectively *càdlàg* if for $\mathbb{P}$ -a.e. $\omega \in \Omega$, the *sample paths* $t \mapsto X_t(\omega) \in \Phi'_\beta$ of X are continuous (respectively right-continuous with left limits).

Let $X = \{X_t\}_{t \in J}$ be a Φ'_β -valued stochastic processes. We say that X is *regular* if for every $t \in J$, X_t is a regular random variable. Some useful properties of Φ'_β -valued regular processes are given below.

Proposition 2.12. *Let $X = \{X_t\}_{t \in J}$ and $Y = \{Y_t\}_{t \in J}$ be Φ'_β -valued regular stochastic processes such that for each $\phi \in \Phi$, $X[\phi] = \{X_t[\phi]\}_{t \in J}$ is a version of $Y = \{Y_t[\phi]\}_{t \in J}$. Then X is a version of Y . Furthermore, if X and Y are right-continuous then they are indistinguishable processes.*

Proof. Fix $t \in J$. Then, as for each $\phi \in \Phi$, $X_t[\phi] = Y_t[\phi]$ $\mathbb{P}$ -a.e., then Proposition 2.11 shows that $X_t = Y_t$ $\mathbb{P}$ -a.e. Therefore, X is a version of Y . Now, assume that both X and Y are right-continuous. Let Ω_X and Ω_Y denote respectively the sets of all $\omega \in \Omega$ such that the maps $t \mapsto X_t(\omega)$ and $t \mapsto Y_t(\omega)$ are right-continuous. Let $\Gamma_{X,Y} = \{\omega \in \Omega : X_t(\omega) = Y_t(\omega), \forall t \in \mathbb{Q}_+\}$, where $\mathbb{Q}_+ = \mathbb{Q} \cap J$. Then, $\mathbb{P}(\Omega_X \cap \Omega_Y \cap \Gamma_{X,Y}) = 1$ and by the right-continuity of X and Y and the denseness of $\mathbb{Q}_+$ in J , it follows by a standard argument that $X_t(\omega) = Y_t(\omega)$ for all $t \in J$, for each $\omega \in \Omega_X \cap \Omega_Y \cap \Gamma_{X,Y}$. Thus, X and Y are indistinguishable. $\square$

3 The Regularization Theorem for Cylindrical Stochastic Processes

Assumption 3.1. *From now on, Φ will always denote a nuclear space over $\mathbb{R}$.*

The main result of this paper is the following theorem that establish conditions for the existence of a Φ'_β -valued, regular, continuous (or càdlàg) version for a cylindrical process in Φ' .

Theorem 3.2 (Regularization Theorem). *Let $X = \{X_t\}_{t \geq 0}$ be a cylindrical process in Φ' satisfying:*

- (1) *For each $\phi \in \Phi$, the real-valued process $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$ has a continuous (respectively càdlàg) version.*
- (2) *For every $T > 0$, the family $\{X_t : t \in [0, T]\}$ of linear maps from Φ into $L^0(\Omega, \mathcal{F}, \mathbb{P})$ is equicontinuous.*

Then, there exists a countably Hilbertian topology ϑ_X on Φ and a $(\widetilde{\Phi_{\vartheta_X}})'_\beta$ -valued continuous (respectively càdlàg) process $Y = \{Y_t\}_{t \geq 0}$, such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \geq 0}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$. Moreover, Y is a Φ'_β -valued, regular, continuous (respectively càdlàg) version of X that is unique up to indistinguishable versions.

We proceed to prove Theorem 3.2. Let $X = \{X_t\}_{t \in [0, T]}$ be a cylindrical process in Φ' satisfying conditions (1) and (2) of Theorem 3.2. Without loss of generality we assume that each $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$ has a continuous version. The càdlàg version case follows from the same arguments.

The next proposition constitutes the main step in the proof of Theorem 3.2.

Proposition 3.3. *For every $T > 0$ there exists a countably Hilbertian topology ϑ_T on Φ and a $(\widetilde{\Phi_{\vartheta_T}})'_\beta$ -valued continuous (respectively càdlàg) process $Y^{(T)} = \{Y_t^{(T)}\}_{t \in [0, T]}$, such that for every $\phi \in \Phi$, $\{Y_t^{(T)}[\phi]\}_{t \in [0, T]}$ is a version of $\{X_t(\phi)\}_{t \in [0, T]}$.*

For the benefit of the reader we split the proof of Proposition 3.3 in several steps. Fix $T > 0$. For each $\phi \in \Phi$, let $\widehat{X}(\phi) = \{\widehat{X}_t(\phi)\}_{t \in [0, T]}$ be a continuous version of $X(\phi) = \{X_t(\phi)\}_{t \in [0, T]}$. It is clear from the corresponding properties of X that $\widehat{X}$ determines a cylindrical process $\widehat{X} = \{\widehat{X}_t\}_{t \in [0, T]}$ in Φ' . Moreover, we have the following result.

Lemma 3.4. *There exists a weaker countably Hilbertian topology θ on Φ such that the family of linear maps $\{\widehat{X}_t : t \in [0, T]\}$ from Φ into $L^0(\Omega, \mathcal{F}, \mathbb{P})$ is θ -equicontinuous.*

Proof. Let $\{\epsilon_n\}_{n \in \mathbb{N}}$ be a decreasing sequence of positive numbers converging to zero. From the equicontinuity of the family of linear maps $\{X_t : t \in [0, T]\}$ and from the fact that $\widehat{X}(\phi)$ is a version of $X(\phi)$ for each $\phi \in \Phi$, it follows that for every $n \in \mathbb{N}$ there exists a continuous Hilbertian semi-norm p_n on Φ such that

$$\mathbb{P}\left(\omega : \left|\widehat{X}_t(\phi)(\omega)\right| > \epsilon_n\right) \leq \epsilon_n, \quad \forall \phi \in B_{p_n}(1), \forall t \in [0, T]. \quad (3.1)$$

Then, if θ is the countably Hilbertian topology on Φ generated by the semi-norms $\{p_n\}_{n \in \mathbb{N}}$, it follows from (3.1) and the fact that $\lim_{n \rightarrow \infty} \epsilon_n = 0$ that the family $\{\widehat{X}_t : t \in [0, T]\}$ is θ -equicontinuous. $\square$

For the next result we need the following terminology. Let $C_T(\mathbb{R})$ and $D_T(\mathbb{R})$ denote respectively the space of continuous and càdlàg real-valued processes defined in $[0, T]$, both are considered equipped with the topology of uniform convergence in probability on $[0, T]$.

Lemma 3.5. *The linear map $\widehat{X}$ from Φ into $C_T(\mathbb{R})$ given by $\phi \mapsto \widehat{X}(\phi) = \{\widehat{X}_t(\phi)\}_{t \in [0, T]}$ is continuous.*

Proof. Let θ be a countably Hilbertian topology on Φ as in Lemma 3.4. Then, for every $t \in [0, T]$, the map $\widehat{X}_t : \Phi_\theta \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is linear and continuous. Therefore, for all $t \in [0, T]$ there exists a continuous and linear map $\widetilde{X}_t : \widetilde{\Phi}_\theta \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ satisfying $\widetilde{X}_t = \widehat{X}_t \circ J$, where J is the canonical embedding from Φ_θ into its completion $\widetilde{\Phi}_\theta$ (see [15], Theorem 2, Section 3.4, p.61-2).

We are going to show that the linear map $\widetilde{X}$ from $\widetilde{\Phi}_\theta$ into $C_T(\mathbb{R})$ given by $\phi \mapsto \widetilde{X}(\phi) = \{\widetilde{X}_t(\phi)\}_{t \in [0, T]}$ is closed. Let $\{\phi_n\}_{n \in \mathbb{N}}$ be a sequence converging to ϕ in $\widetilde{\Phi}_\theta$ and assume that there exists $Y \in C_T(\mathbb{R})$ such that $\sup_{t \in [0, T]} |\widetilde{X}_t(\phi_n) - Y_t|$ converges in probability to 0 as $n \rightarrow \infty$. We have to prove that $\widetilde{X}(\phi) = Y$ in $C_T(\mathbb{R})$.

First, for every $t \in [0, T]$ the continuity of the map $\widetilde{X}_t : \widetilde{\Phi}_\theta \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ and the fact that $\{\phi_n\}_{n \in \mathbb{N}}$ converges to ϕ in $\widetilde{\Phi}_\theta$ implies that the sequence of random variables $\{\widetilde{X}_t(\phi_n)\}_{n \in \mathbb{N}}$ converges in probability to $\widetilde{X}_t(\phi)$.

On the other hand, as $\sup_{t \in [0, T]} |\widetilde{X}_t(\phi_n) - Y_t|$ converges in probability to 0 as $n \rightarrow \infty$, then for every $t \in [0, T]$ the sequence of random variables $\{\widetilde{X}_t(\phi_n)\}_{n \in \mathbb{N}}$ converges in probability to Y_t . Therefore, by uniqueness of limits in $L^0(\Omega, \mathcal{F}, \mathbb{P})$ it follows that $\widetilde{X}_t(\phi) = Y_t$ $\mathbb{P}$ -a.e. for every $t \in [0, T]$, and hence $\widetilde{X}(\phi) = Y$ in $C_T(\mathbb{R})$.

Then, the linear map $\widetilde{X} : \widetilde{\Phi}_\theta \rightarrow C_T(\mathbb{R})$ is sequentially closed and therefore closed because $\widetilde{\Phi}_\theta$ is pseudo-metrizable (Proposition 2.4) and hence first-countable. Now, because $\widetilde{\Phi}_\theta$ is a Baire space (Proposition 2.4) and $C_T(\mathbb{R})$ is a complete, metrizable, topological vector space, the closed graph theorem (see [26], Theorem 14.3.4, p.465) shows that $\widetilde{X}$ is continuous.

Now, because for every $t \in [0, T]$, $\widehat{X}_t = \widetilde{X}_t \circ J$ then we have that the map $\widehat{X}$ from Φ_θ into $C_T(\mathbb{R})$ given by $\phi \mapsto \widehat{X}(\phi) = \{\widehat{X}_t(\phi)\}_{t \in [0, T]}$ satisfies $\widehat{X} = \widetilde{X} \circ J$. Therefore, as both $\widetilde{X}$ and J are continuous, it follows that $\widehat{X}$ is also continuous.

Moreover, as the topology θ is weaker than the nuclear topology on Φ , it follows that $\widehat{X}$ is continuous as a map from Φ into $C_T(\mathbb{R})$. $\square$

Remark 3.6. *In the case that for each $\phi \in \Phi$, $X(\phi) = \{X_t(\phi)\}_{t \in [0, T]}$ has a càdlàg version $\widehat{X}(\phi) = \{\widehat{X}_t(\phi)\}_{t \in [0, T]}$, then in Lemma 3.5 we have that the linear mapping $\widehat{X}$ from Φ into $D_T(\mathbb{R})$ given by $\phi \mapsto \widehat{X}(\phi)$ is continuous.*

Lemma 3.7. *Let D be a countable dense subset of $[0, T]$. For every $\epsilon > 0$ there exists a continuous Hilbertian semi-norm p on Φ such that*

$$\mathbb{E} \left(\sup_{t \in D} |1 - e^{i\widehat{X}_t(\phi)}| \right) \leq \epsilon + 2p(\phi)^2, \quad \forall \phi \in \Phi. \quad (3.2)$$

Proof. We proceed in a way similar to Lemma 1 of Mitoma [24]. Let $\epsilon > 0$. From the continuity of the exponential function there exists $\delta > 0$ such that $|1 - e^{ir}| \leq \frac{\epsilon}{2}$ if $|r| \leq \delta$. Now, from the continuity of the map $\phi \mapsto \widehat{X}(\phi)$ from Φ into $C_T(\mathbb{R})$ (Lemma 3.5), there exists a continuous Hilbertian semi-norm p on Φ such that

$$\mathbb{P} \left(\omega : \sup_{t \in D} |\widehat{X}_t(\phi)(\omega)| > \delta \right) \leq \frac{\epsilon}{4}, \quad \forall \phi \in B_p(1). \quad (3.3)$$

Let $\Upsilon = \{\omega \in \Omega : \sup_{t \in D} |\widehat{X}_t(\phi)(\omega)| < \delta\}$. Then, if $\phi \in B_p(1)$ it follows from (3.3) that

$$\mathbb{E} \left(\sup_{t \in D} |1 - e^{i\widehat{X}_t(\phi)}| \right) \leq \int_{\Upsilon} \sup_{t \in D} |1 - e^{i\widehat{X}_t(\phi)(\omega)}| \mathbb{P}(d\omega) + 2\mathbb{P}(\Upsilon^c) \leq \epsilon.$$

On the other hand, because $\sup_{t \in D} |1 - e^{i\widehat{X}_t(\phi)}| \leq 2$ for any $\phi \in \Phi$, then if $\phi \in B_p(1)^c$, we have

$$\mathbb{E} \left(\sup_{t \in D} |1 - e^{i\widehat{X}_t(\phi)}| \right) \leq 2p(\phi)^2.$$

Therefore, from the above inequalities we obtain (3.2). $\square$

Lemma 3.8. *There exists an increasing sequence of continuous Hilbertian semi-norms $\{q_n\}_{n \in \mathbb{N}}$ on Φ , a subset Ω_Y of Ω such that $\mathbb{P}(\Omega_Y) = 1$ and a sequence of stochastic processes $Y^{(n)} = \{Y_t^{(n)}\}_{t \in [0, T]}$, $n \in \mathbb{N}$, satisfying:*

- (1) *For each $n \in \mathbb{N}$, $Y^{(n)}$ is a Φ'_{q_n} -valued process such that for every $\phi \in \Phi_{q_n}$, $Y^{(n)}[\phi] = \{Y_t^{(n)}[\phi]\}_{t \in [0, T]}$ is a continuous real-valued process.*
- (2) *For each $\omega \in \Omega_Y$, there exists $N(\omega)$ such that*
 - (a) *For all $n \geq N(\omega)$, $\sup_{t \in [0, T]} q'_n(Y_t^{(n)}(\omega)) < \infty$, and*
 - (b) *For all $m \geq n \geq N(\omega)$, $Y_t^{(m)}(\omega) = i'_{q_n, q_m} Y_t^{(n)}(\omega)$ for all $t \in [0, T]$.*
- (3) *For every $\phi \in \Phi$, there exists $\Delta_\phi \subseteq \Omega$ with $\mathbb{P}(\Delta_\phi) = 1$ such that for every $\omega \in \Omega_Y \cap \Delta_\phi$ there exists $N(\omega)$ such that for each $n \geq N(\omega)$, $Y_t^{(n)}(\omega)[i_{q_n} \phi] = \widehat{X}_t(\phi)(\omega)$ for all $t \in [0, T]$.*

Proof. We follow similar arguments to those used by Itô and Nawata in [14]. Let D be a countable dense subset of $[0, T]$. Let $\{\epsilon_n\}_{n \in \mathbb{N}}$ be a sequence of positive numbers such that $\sum_{n \in \mathbb{N}} \epsilon_n < \infty$. From Lemma 3.7 there exist an increasing sequence of continuous Hilbertian semi-norms $\{p_n\}_{n \in \mathbb{N}}$ on Φ such that for each $n \in \mathbb{N}$, ϵ_n and p_n satisfy (3.2).

Now, as Φ is nuclear, there exists an increasing sequence of continuous Hilbertian semi-norms $\{q_n\}_{n \in \mathbb{N}}$ on Φ such that for each $n \in \mathbb{N}$, $p_n \leq q_n$ and the inclusion i_{p_n, q_n} is Hilbert-Schmidt. Let α be the countably Hilbertian topology on Φ generated by the semi-norms $\{q_n\}_{n \in \mathbb{N}}$. The space Φ_α is separable (Proposition 2.4). Let $B = \{\xi_k : k \in \mathbb{N}\}$ be a countable dense subset of Φ_α . For every $n \in \mathbb{N}$, from

an application of the Schmidt orthogonalization procedure to B , we can find a complete orthonormal system $\{\phi_j^{q_n}\}_{j \in \mathbb{N}} \subseteq \Phi$ of Φ_{q_n} , such that

$$\xi_k = \sum_{j=1}^k a_{j,k,n} \phi_j^{q_n} + \varphi_{k,n}, \quad \forall k \in \mathbb{N}, \quad (3.4)$$

with $a_{j,k,n} \in \mathbb{R}$ and $\varphi_{k,n} \in \text{Ker}(q_n)$, for each $j, k \in \mathbb{N}$.

Let $n \in \mathbb{N}$. From the inequality: $1 - e^{-r/2} \geq 1 - e^{-1/2} = \frac{\sqrt{e}-1}{\sqrt{e}}$ for $r > 1$, for any $C > 0$ we have

$$\begin{aligned} & \mathbb{P} \left(\sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\phi_j^{q_n}) \right|^2 > C^2 \right) \\ & \leq \frac{\sqrt{e}}{\sqrt{e}-1} \mathbb{E} \left(1 - \exp \left\{ -\frac{1}{2C^2} \sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\phi_j^{q_n}) \right|^2 \right\} \right) \\ & = \lim_{m \rightarrow \infty} \frac{\sqrt{e}}{\sqrt{e}-1} \mathbb{E} \sup_{t \in D} \left(1 - \exp \left\{ -\frac{1}{2C^2} \sum_{j=1}^m \left| \widehat{X}_t(\phi_j^{q_n}) \right|^2 \right\} \right) \end{aligned} \quad (3.5)$$

Now, setting $\phi = \sum_{j=1}^m z_j \phi_j^{q_n}$ for $z_1, \dots, z_m \in \mathbb{R}$, in (3.2) for p_n we have

$$\mathbb{E} \left(\sup_{t \in D} \left| 1 - \exp \left\{ i \sum_{j=1}^m z_j \widehat{X}_t(\phi_j^{q_n}) \right\} \right| \right) \leq \epsilon_n + 2 \sum_{j=1}^m z_j^2 p_n(\phi_j^{q_n})^2. \quad (3.6)$$

Integrating both sides of (3.6) with respect to $\prod_{j=1}^m N_C(dz_j)$, where N_C is the centered Gaussian measure on $\mathbb{R}$ with variance $1/C^2$, we have

$$\int_{\mathbb{R}^m} \mathbb{E} \left(\sup_{t \in D} \left| 1 - \exp \left\{ i \sum_{j=1}^m z_j \widehat{X}_t(\phi_j^{q_n}) \right\} \right| \right) \prod_{j=1}^m N_C(dz_j) \leq \epsilon_n + \frac{2}{C^2} \sum_{j=1}^m p_n(\phi_j^{q_n})^2. \quad (3.7)$$

On the other hand, as $\prod_{j=1}^m N_C(dz_j)$ is a Gaussian measure on $\mathbb{R}^m$, for each $t \in [0, T]$ and $\omega \in \Omega$ we have

$$\exp \left\{ -\frac{1}{2C^2} \sum_{j=1}^m \left| \widehat{X}_t(\phi_j^{q_n})(\omega) \right|^2 \right\} = \int_{\mathbb{R}^m} \exp \left\{ i \sum_{j=1}^m z_j \widehat{X}_t(\phi_j^{q_n})(\omega) \right\} \prod_{j=1}^m N_C(dz_j), \quad (3.8)$$

and therefore from (3.8) and the Fubini theorem it follows that

$$\begin{aligned} & \mathbb{E} \sup_{t \in D} \left(1 - \exp \left\{ -\frac{1}{2C^2} \sum_{j=1}^m \left| \widehat{X}_t(\phi_j^{q_n}) \right|^2 \right\} \right) \\ & \leq \int_{\mathbb{R}^m} \mathbb{E} \left(\sup_{t \in D} \left| 1 - \exp \left\{ i \sum_{j=1}^m z_j \widehat{X}_t(\phi_j^{q_n}) \right\} \right| \right) \prod_{j=1}^m N_C(dz_j). \end{aligned} \quad (3.9)$$

Then, from (3.5), (3.7) and (3.9), it follows that

$$\begin{aligned} \mathbb{P} \left(\sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\phi_j^{q_n}) \right|^2 > C^2 \right) & \leq \lim_{m \rightarrow \infty} \frac{\sqrt{e}}{\sqrt{e}-1} \left(\epsilon_n + \frac{2}{C^2} \sum_{j=1}^m p_n(\phi_j^{q_n})^2 \right) \\ & = \frac{\sqrt{e}}{\sqrt{e}-1} \left(\epsilon_n + \frac{2}{C^2} \|i_{p_n, q_n}\|_{\mathcal{L}_2(\Phi_{q_n}, \Phi_{p_n})}^2 \right), \end{aligned}$$

where $\|i_{p_n, q_n}\|_{\mathcal{L}_2(\Phi_{q_n}, \Phi_{p_n})} < \infty$ as i_{p_n, q_n} is Hilbert-Schmidt. Letting $C \rightarrow \infty$, we get

$$\mathbb{P} \left(\sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\phi_j^{q_n}) \right|^2 < \infty \right) \geq 1 - \frac{\sqrt{e}}{\sqrt{e}-1} \epsilon_n. \quad (3.10)$$

Following the same arguments as above but now replacing $\phi_j^{q_n}$ for $\varphi_{j,n}$ and using the fact that $\varphi_{j,n} \in \text{Ker}(p_n)$ for each $j \in \mathbb{N}$ (recall $p_n \leq q_n$), we have that

$$\mathbb{P} \left(\sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\varphi_{j,n}) \right|^2 > 0 \right) \leq \frac{\sqrt{e}}{\sqrt{e}-1} \epsilon_n. \quad (3.11)$$

Then, if we define $\Omega_n \subseteq \Omega$ by

$$\Omega_n = \left\{ \omega : \sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\phi_j^{q_n})(\omega) \right|^2 < \infty \text{ and } \widehat{X}_t(\varphi_{j,n})(\omega) = 0, \forall t \in D, j \in \mathbb{N} \right\}, \quad (3.12)$$

it follows from (3.10) and (3.11) that

$$\mathbb{P}(\Omega_n) \geq 1 - 2 \frac{\sqrt{e}}{\sqrt{e}-1} \epsilon_n. \quad (3.13)$$

The next point in our agenda is to define the set Ω_Y of $\mathbb{P}$ -measure 1 and the stochastic processes $Y^{(n)} = \{Y_t^{(n)}\}_{t \in [0, T]}$, $n \in \mathbb{N}$, that satisfy the properties (1)-(3) in the statement of the Lemma. But before, we set some additional notation.

For every $n \in \mathbb{N}$, let $\Gamma_n \subseteq \Omega$ given by

$$\Gamma_n = \left\{ \omega : \forall j \in \mathbb{N}, t \mapsto \widehat{X}_t(\phi_j^{q_n})(\omega) \text{ is continuous} \right\}. \quad (3.14)$$

Then, for each $n \in \mathbb{N}$ we have $\mathbb{P}(\Gamma_n) = 1$. Also, for each $n \in \mathbb{N}$ define $A_n \subseteq \Omega$ by

$$A_n = \left\{ \omega : \widehat{X}_t(\xi_k)(\omega) = \sum_{j=1}^k a_{j,k,n} \widehat{X}_t(\phi_j^{q_n})(\omega) + \widehat{X}_t(\varphi_{k,n})(\omega), \forall k \in \mathbb{N}, t \in D \right\}. \quad (3.15)$$

For every $n \in \mathbb{N}$, it follows from (3.4) and the linearity of each $\widehat{X}_t$ that $\mathbb{P}(A_n) = 1$. Now, for $n \in \mathbb{N}$ define

$$\Lambda_n = \Omega_n \cap \Gamma_n \cap A_n. \quad (3.16)$$

Then, it follows from (3.13) and the fact that $\mathbb{P}(\Gamma_n) = 1$ and $\mathbb{P}(A_n) = 1$ that

$$\mathbb{P}(\Lambda_n) \geq 1 - 2 \frac{\sqrt{e}}{\sqrt{e}-1} \epsilon_n. \quad (3.17)$$

We are ready to define the stochastic processes $Y^{(n)}$, $n \in \mathbb{N}$. For each $n \in \mathbb{N}$, let $\{f_j^{q_n}\}_{j \in \mathbb{N}}$ be a complete orthonormal system in Φ'_{q_n} dual to $\{\phi_j^{q_n}\}_{j \in \mathbb{N}}$, i.e. $f_j^{q_n}[\phi_i^{q_n}] = \delta_{i,j}$ where $\delta_{i,j} = 1$ if $i = j$ and $\delta_{i,j} = 0$ if $i \neq j$. For each $t \in [0, T]$, we define

$$Y_t^{(n)}(\omega) := \begin{cases} \sum_{j=1}^{\infty} \widehat{X}_t(\phi_j^{q_n})(\omega) f_j^{q_n}, & \text{for } \omega \in \Lambda_n, \\ 0, & \text{elsewhere.} \end{cases} \quad (3.18)$$

Note that $Y^{(n)} = \{Y_t^{(n)}\}_{t \in [0, T]}$ is a well-defined Φ'_{q_n} -valued stochastic process. This is because if $\omega \in \Lambda_n$, then the infinite sum in (3.18) is convergent, as from Parseval's identity, (3.12), (3.16) and (3.18), we have:

$$\sup_{t \in [0, T]} q'_n(Y_t^{(n)}(\omega))^2 = \sup_{t \in D} \sum_{j=1}^{\infty} \left| \widehat{X}_t(\phi_j^{q_n})(\omega) \right|^2 < \infty. \quad (3.19)$$

Moreover, it follows from (3.14), (3.16) and (3.18) that for every $\phi \in \Phi_{q_n}$, $Y^{(n)}[\phi]$ is a continuous real-valued process. Therefore, $Y^{(n)}$ satisfies the property (1) in the statement of the Lemma.

Also, from (3.18) it follows that for each $\omega \in \Lambda_n$, $Y_t^{(n)}(\omega)[i_{q_n}\phi_j^{q_n}] = \widehat{X}_t(\phi_j^{q_n})(\omega)$, for all $j \in \mathbb{N}$ and $t \in [0, T]$. Similarly, from the fact that $q_n(\varphi_{j,n}) = 0$ for all $j \in \mathbb{N}$, we have $|f_j^{q_n}[i_{q_n}\varphi_{j,n}]| \leq q'_n(f_j^{q_n})q_n(i_{q_n}\varphi_{j,n}) = 0$ for all $j \in \mathbb{N}$, then (3.12), (3.16) and (3.18) implies that for each $\omega \in \Lambda_n$, $Y_t^{(n)}(\omega)[i_{q_n}\varphi_{j,n}] = \widehat{X}_t(\varphi_{j,n})(\omega) = 0$ for all $t \in D$, $j \in \mathbb{N}$. So, by (3.14), (3.15) and (3.16) we have

$$\forall \omega \in \Lambda_n, \quad Y_t^{(n)}(\omega)[i_{q_n}\xi_k] = \widehat{X}_t(\xi_k)(\omega), \quad \forall k \in \mathbb{N}, t \in [0, T]. \quad (3.20)$$

Now we are going to show that (3.20) implies that for every $\phi \in \Phi$, $Y_t^{(n)}[i_{q_n}\phi] = \widehat{X}_t(\phi)$ $\mathbb{P}$ -a.e. on Λ_n , for all $t \in [0, T]$.

Let $\phi \in \Phi$. Since $B = \{\xi_k : k \in \mathbb{N}\}$ is dense in Φ_α , there exists a sequence $\{\xi_{k_j}\}_{j \in \mathbb{N}} \subseteq B$ that α -converges to ϕ . As α is the countably Hilbertian topology generated by the semi-norms $\{q_n\}_{n \in \mathbb{N}}$, then $\{i_{q_n}\xi_{k_j}\}_{j \in \mathbb{N}}$ converges to $i_{q_n}\phi$ in Φ_{q_n} as $j \rightarrow \infty$. Therefore, $Y_t^{(n)}(\omega)[i_{q_n}\xi_{k_j}] \rightarrow Y_t^{(n)}(\omega)[i_{q_n}\phi]$ as $j \rightarrow \infty$, for all $t \in [0, T]$ and $\omega \in \Omega$.

On the other hand, observe that (3.3) implies that $\widehat{X} : \Phi \rightarrow C_T(\mathbb{R})$, $\phi \mapsto \widehat{X}(\phi)$, is continuous with respect to the countably Hilbertian topology on Φ generated by the semi-norms $\{p_n\}_{n \in \mathbb{N}}$ and since this topology is weaker than α (because $p_n \leq q_n$ for all $n \in \mathbb{N}$), then it is also α -continuous. Therefore, there exists $\Delta_\phi \subseteq \Omega$ with $\mathbb{P}(\Delta_\phi) = 1$ and a subsequence $\{\xi_{k_{j,\nu}}\}_{\nu \in \mathbb{N}}$ of $\{\xi_{k_j}\}_{j \in \mathbb{N}}$ such that for all $\omega \in \Delta_\phi$, $\widehat{X}_t(\xi_{k_{j,\nu}})(\omega) \rightarrow \widehat{X}_t(\phi)(\omega)$, as $\nu \rightarrow \infty$, for all $t \in [0, T]$.

Then, (3.20) and the uniqueness of limits implies that

$$\forall \omega \in \Lambda_n \cap \Delta_\phi, \quad Y_t^{(n)}(\omega)[i_{q_n}\phi] = \widehat{X}_t(\phi)(\omega), \quad \forall t \in [0, T]. \quad (3.21)$$

Our final step is to define the set Ω_Y and to verify that it and the processes $Y^{(n)}$, $n \in \mathbb{N}$, defined in (3.18) satisfy the conditions (2) and (3) of the statement of the Lemma. First, it follows from (3.17), our assumption that $\sum_{n \in \mathbb{N}} \epsilon_n < \infty$, and the Borel-Cantelli lemma that

$$\mathbb{P}(\Omega_Y) = 1, \quad \text{where} \quad \Omega_Y := \bigcup_{N \in \mathbb{N}} \bigcap_{n \geq N} \Lambda_n. \quad (3.22)$$

Let $\omega \in \Omega_Y$. Then, it follows from (3.19) and (3.22) that there exists some $N(\omega)$ such that for all $n \geq N(\omega)$, $\sup_{t \in D} q'_n(Y_t^{(n)}(\omega)) < \infty$. Thus, the property (2)(a) in the statement of the Lemma is satisfied. Moreover, from (3.20) and (3.22) there exists $N(\omega)$ such that for all $m \geq n \geq N(\omega)$, for every $k \in \mathbb{N}$ and $t \in [0, T]$ we have $Y_t^{(m)}(\omega)[i_{q_m}\xi_k] = Y_t^{(n)}(\omega)[i_{q_n}\xi_k]$. But as $B = \{\xi_j\}_{j \in \mathbb{N}}$ is dense in Φ_α , and therefore in Φ_{q_m} and in Φ_{q_n} , then it follows that for all $t \in [0, T]$, $Y_t^{(m)}(\omega)[i_{q_m}\phi] = Y_t^{(n)}(\omega)[i_{q_n}\phi]$ for all $\phi \in \Phi$, that is $Y_t^{(m)}(\omega) = i'_{q_n, q_m} Y_t^{(n)}(\omega)$. Hence, the property (2)(b) of the statement of the Lemma is also satisfied.

Finally, the property (3) of the statement of the Lemma is a consequence of (3.21) and (3.22). $\square$

Proof of Proposition 3.3. We use similar arguments to those used by Mitoma in [24]. Let $\{q_n\}_{n \in \mathbb{N}}$, Ω_Y and $\{Y^{(n)}\}_{n \in \mathbb{N}}$ be as given in Lemma 3.8.

Let $\{\varrho_n\}_{n \in \mathbb{N}}$ be an increasing sequence of continuous Hilbertian semi-norm on Φ such that for every $n \in \mathbb{N}$, $q_n \leq \varrho_n$ and i_{q_n, ϱ_n} is Hilbert-Schmidt. Let ϑ be the countably Hilbertian topology on Φ generated by the semi-norms $\{\varrho_n\}_{n \in \mathbb{N}}$. The topology ϑ is weaker than the nuclear topology on Φ .

Moreover, from Proposition 2.4 we have that the completion $\widetilde{\Phi}_\vartheta$ of Φ_ϑ satisfies

$$(\widetilde{\Phi}_\vartheta)' = \bigcup_{n \in \mathbb{N}} \Phi'_{\varrho_n}. \quad (3.23)$$

Now, for every $n \in \mathbb{N}$ the map i_{ϱ_n} is linear and continuous from Φ_ϑ into Φ_{ϱ_n} , and therefore i_{ϱ_n} has a unique continuous and linear extension $\widetilde{i}_{\varrho_n} : \widetilde{\Phi}_\vartheta \rightarrow \Phi_{\varrho_n}$ (see [15], Theorem 2, Section 3.4, p.61-2). Hence, for each $n \in \mathbb{N}$ the dual map $(\widetilde{i}_{\varrho_n})' : \Phi'_{\varrho_n} \rightarrow (\widetilde{\Phi}_\vartheta)'_\beta$ is linear and continuous, and corresponds to the canonical inclusion from Φ'_{ϱ_n} into $(\widetilde{\Phi}_\vartheta)'_\beta$.

Let $Y = \{Y_t\}_{t \in [0, T]}$ be defined for each $t \in [0, T]$ by

$$Y_t(\omega) := \begin{cases} i'_{q_n, \varrho_n} Y_t^{(n)}(\omega), & \text{for } \omega \in \Omega_Y, n \geq N(\omega), \\ 0, & \text{elsewhere.} \end{cases} \quad (3.24)$$

For every $t \in [0, T]$, Y_t is well-defined $(\widetilde{\Phi}_\vartheta)'_\beta$ -valued random variable. In effect, it is clear from (3.23) and (3.24) that Y_t takes values in $(\widetilde{\Phi}_\vartheta)'$. Now, as for all $m, n \in \mathbb{N}$, $m \geq n$, we have $q_n \leq q_m$, $q_n \leq \varrho_n$, and $q_m \leq \varrho_m$, then the following diagram commutes:

$$\begin{array}{ccc} \Phi'_{q_n} & \xrightarrow{i'_{q_n, \varrho_n}} & \Phi'_{\varrho_n} \\ i'_{q_n, q_m} \downarrow & & \downarrow i'_{\varrho_n, \varrho_m} \\ \Phi'_{q_m} & \xrightarrow{i'_{q_m, \varrho_m}} & \Phi'_{\varrho_m} \end{array} \quad (3.25)$$

Then Lemma 3.8(2)(b) and (3.25) implies that for each $\omega \in \Omega_Y$, if $m \geq n \geq N(\omega)$ we have

$$i'_{q_m, \varrho_m} Y_t^{(m)}(\omega) = i'_{q_m, \varrho_m} \circ i'_{q_n, q_m} Y_t^{(n)}(\omega) = i'_{\varrho_n, \varrho_m} \circ i'_{q_n, \varrho_n} Y_t^{(n)}(\omega), \quad \forall t \in [0, T].$$

Therefore, Y_t is well-defined. Finally, the fact that Y_t is $\mathcal{B}((\widetilde{\Phi}_\vartheta)'_\beta)/\mathcal{F}$ -measurable map is a consequence of (3.24), the fact that $Y_t^{(n)}$ is a Φ'_{q_n} -valued random variable and that i'_{q_n, ϱ_n} is continuous, for every $n \in \mathbb{N}$.

Now we are going to show that Y is a $(\widetilde{\Phi}_\vartheta)'_\beta$ -valued continuous process. For every $n \in \mathbb{N}$, let $\{\phi_j^{\varrho_n}\}_{j \in \mathbb{N}} \subseteq \Phi$ be a complete orthonormal system in Φ_{ϱ_n} . Fix $\omega \in \Omega_Y$ and let $n \geq N(\omega)$. Then, from the definition of the dual operator i'_{q_n, ϱ_n} of i_{q_n, ϱ_n} and Lemma 3.8(1)-(2)(a), we have

$$\begin{aligned} \sum_{j=1}^{\infty} \sup_{t \in [0, T]} \left| i'_{q_n, \varrho_n} Y_t^{(n)}(\omega) [\phi_j^{\varrho_n}] \right|^2 &\leq \sum_{j=1}^{\infty} \sup_{t \in [0, T]} q'_n(Y_t^{(n)}(\omega))^2 q_n(i_{q_n, \varrho_n} \phi_j^{\varrho_n})^2 \\ &= \left(\sup_{t \in [0, T]} q'_n(Y_t^{(n)}(\omega))^2 \right) \|i_{q_n, \varrho_n}\|_{\mathcal{L}_2(\Phi_{\varrho_n}, \Phi_{q_n})}^2 \\ &< \infty, \end{aligned} \quad (3.26)$$

where $\|i_{q_n, \varrho_n}\|_{\mathcal{L}_2(\Phi_{\varrho_n}, \Phi_{q_n})} < \infty$ because i_{q_n, ϱ_n} is Hilbert-Schmidt.

Next we prove the right continuity of the map $t \mapsto Y_t(\omega)$ in Φ'_{ϱ_n} . Let $0 \leq t < T$. Then, from (3.24), Parseval's identity, (3.26), the dominated convergence theorem

and the continuity of each map $t \mapsto Y_t^{(n)}(\omega)[i_{q_n, \varrho_n} \phi_j^{\varrho_n}]$ (Lemma 3.8(1)), we have

$$\begin{aligned} \lim_{s \rightarrow t+} \varrho'_n(Y_t(\omega) - Y_s(\omega))^2 &= \lim_{s \rightarrow t+} \sum_{j=1}^{\infty} \left| i'_{q_n, \varrho_n}(Y_t^{(n)}(\omega) - Y_s^{(n)}(\omega))[\phi_j^{\varrho_n}] \right|^2 \\ &= \sum_{j=1}^{\infty} \lim_{s \rightarrow t+} \left| Y_t^{(n)}(\omega)[i_{q_n, \varrho_n} \phi_j^{\varrho_n}] - Y_s^{(n)}(\omega)[i_{q_n, \varrho_n} \phi_j^{\varrho_n}] \right|^2 \\ &= 0. \end{aligned}$$

Therefore, the map $t \mapsto Y_t(\omega)$ is right-continuous in Φ'_{ϱ_n} . The left continuity of $t \mapsto Y_t(\omega)$ in Φ'_{ϱ_n} can be proven similarly. Moreover, as the canonical inclusion $(\tilde{i}_{\varrho_n})' : \Phi'_{\varrho_n} \rightarrow (\widetilde{\Phi_{\vartheta}})'_{\beta}$ is continuous, then the continuity of $t \mapsto Y_t(\omega)$ in Φ'_{ϱ_n} implies that the map $t \mapsto (\tilde{i}_{\varrho_n})' Y_t(\omega)$ is continuous from $[0, \infty)$ into $(\widetilde{\Phi_{\vartheta}})'_{\beta}$. Hence, Y is a $(\widetilde{\Phi_{\vartheta}})'_{\beta}$ -valued continuous process.

Finally, the fact that for every $\phi \in \Phi$, $Y[\phi]$ is a version of $X(\phi)$ is a direct consequence of Lemma 3.8(3), (3.24) and because $\hat{X}(\phi)$ is a version of $X(\phi)$. $\square$

Proof of Theorem 3.2. Let $\{T_k\}_{k \in \mathbb{N}}$ an increasing sequence of positive numbers such that $\lim_{k \rightarrow \infty} T_k = \infty$. From Proposition 3.3, for every $k \in \mathbb{N}$ there exists a countably Hilbertian topology ϑ_{T_k} on Φ determined by an increasing sequence $\{\varrho_{k,n}\}_{n \in \mathbb{N}}$ of continuous Hilbertian semi-norms on Φ , and a $(\widetilde{\Phi_{\vartheta_{T_k}}})'_{\beta}$ -valued continuous process $Y^{(T_k)} = \{Y_t^{(T_k)}\}_{t \in [0, T_k]}$, such that for every $\phi \in \Phi$, $\{Y_t^{(T_k)}[\phi]\}_{t \in [0, T_k]}$ is a version of $\{X_t(\phi)\}_{t \in [0, T_k]}$.

Without loss of generality we can assume that for every $k \in \mathbb{N}$, $\varrho_{k,n} \leq \varrho_{k+1,n}$, for all $n \in \mathbb{N}$. This implies that for every $k \in \mathbb{N}$, the topology $\vartheta_{T_{k+1}}$ is weaker than ϑ_{T_k} and therefore that the canonical inclusion $i_{\vartheta_{T_k}, \vartheta_{T_{k+1}}} : \Phi_{\vartheta_{T_{k+1}}} \rightarrow \Phi_{\vartheta_{T_k}}$ is linear and continuous. Then, this map has a unique continuous and linear extension $\tilde{i}_{\vartheta_k, \vartheta_{k+1}} : \widetilde{\Phi_{\vartheta_{T_{k+1}}}} \rightarrow \widetilde{\Phi_{\vartheta_{T_k}}}$. Hence, the dual map $(\tilde{i}_{\vartheta_{T_k}, \vartheta_{T_{k+1}}})' : (\widetilde{\Phi_{\vartheta_{T_k}}})'_{\beta} \rightarrow (\widetilde{\Phi_{\vartheta_{T_{k+1}}}})'_{\beta}$ is also linear and continuous, and corresponds to the canonical inclusion from $(\widetilde{\Phi_{\vartheta_{T_k}}})'_{\beta}$ into $(\widetilde{\Phi_{\vartheta_{T_{k+1}}}})'_{\beta}$.

Let ϑ_X be the countably Hilbertian topology on Φ generated by the semi-norms $\{\varrho_{k,n} : n, k \in \mathbb{N}\}$. Then, for each $k \in \mathbb{N}$ the topology ϑ_{T_k} is weaker than ϑ_X . Hence the canonical inclusion $i_{\vartheta_{T_k}, \vartheta_X}$ from Φ_{ϑ_X} into $\Phi_{\vartheta_{T_k}}$ is linear and continuous. This map has an extension $\tilde{i}_{\vartheta_{T_k}, \vartheta_X} : \widetilde{\Phi_{\vartheta_X}} \rightarrow \widetilde{\Phi_{\vartheta_{T_k}}}$ that is also linear and continuous. The dual operator $(\tilde{i}_{\vartheta_{T_k}, \vartheta_X})' : (\widetilde{\Phi_{\vartheta_{T_k}}})'_{\beta} \rightarrow (\widetilde{\Phi_{\vartheta_X}})'_{\beta}$ corresponds to the canonical inclusion from $(\widetilde{\Phi_{\vartheta_{T_k}}})'_{\beta}$ into $(\widetilde{\Phi_{\vartheta_X}})'_{\beta}$ and is linear and continuous. For every $k \in \mathbb{N}$, one can easily verify that $\tilde{i}_{\vartheta_{T_k}, \vartheta_X} = \tilde{i}_{\vartheta_k, \vartheta_{k+1}} \circ \tilde{i}_{\vartheta_{T_{k+1}}, \vartheta_X}$ and hence $(\tilde{i}_{\vartheta_{T_k}, \vartheta_X})' = (\tilde{i}_{\vartheta_{T_{k+1}}, \vartheta_X})' \circ (\tilde{i}_{\vartheta_k, \vartheta_{k+1}})'$.

Take $Y = \{Y_t\}_{t \geq 0}$ defined by the prescription $Y_t = Y_t^{(T_k)}$ if $t \in [0, T_k]$. Then Y is a $(\widetilde{\Phi_{\vartheta_X}})'_{\beta}$ -valued continuous process. To prove this, note that for any $k \in \mathbb{N}$, $\{(\tilde{i}_{\vartheta_{T_k}, \vartheta_{T_{k+1}}})' Y_t^{(T_k)}\}_{t \in [0, T_k]}$ and $\{Y_t^{(T_{k+1})}\}_{t \in [0, T_k]}$ are continuous $(\widetilde{\Phi_{\vartheta_{T_{k+1}}}})'_{\beta}$ -valued processes. Moreover, for every $\phi \in \Phi$ and $t \in T_k$ we have $\mathbb{P}$ -a.e.

$$\begin{aligned} (\tilde{i}_{\vartheta_{T_k}, \vartheta_{T_{k+1}}})' Y_t^{(T_k)}[\tilde{i}_{\vartheta_{T_{k+1}}, \vartheta_X} \phi] &= Y_t^{(T_k)}[\tilde{i}_{\vartheta_{T_k}, \vartheta_{T_{k+1}}} \circ \tilde{i}_{\vartheta_{T_{k+1}}, \vartheta_X} \phi] \\ &= Y_t^{(T_k)}[\tilde{i}_{\vartheta_{T_k}, \vartheta_X} \phi] = X_t(\phi) = Y_t^{(T_{k+1})}[\tilde{i}_{\vartheta_{T_{k+1}}, \vartheta_X} \phi]. \end{aligned}$$

Then, Proposition 2.12 shows that $\{(\tilde{i}_{\vartheta_{T_k}, \vartheta_{T_{k+1}}})' Y_t^{(T_k)}\}_{t \in [0, T_k]}$ and $\{Y_t^{(T_{k+1})}\}_{t \in [0, T_k]}$

are indistinguishable processes in $(\widetilde{\Phi_{\vartheta_{T_k+1}}})'_\beta$. Therefore, Y is well-defined $(\widetilde{\Phi_{\vartheta_X}})'_\beta$ -valued process.

Now, because for each $k \in \mathbb{N}$ the $(\widetilde{\Phi_{\vartheta_{T_k}}})'_\beta$ -valued process $\{Y_t^{(T_k)}\}_{t \in [0, T_k]}$ is continuous and the map $(\widetilde{i_{\vartheta_{T_k}, \vartheta_X}})'$ is continuous, it follows that $\{Y_t\}_{t \in [0, T_k]} = \{(\widetilde{i_{\vartheta_{T_k}, \vartheta_X}})' Y_t^{(T_k)}\}_{t \in [0, T_k]}$ is a $(\widetilde{\Phi_{\vartheta_X}})'_\beta$ -valued continuous process. Therefore, Y is a continuous processes in $(\widetilde{\Phi_{\vartheta_X}})'_\beta$.

Moreover, by the properties of the processes $Y^{(T_k)}$ and the definition of Y it follows that for every $\phi \in \Phi$, $Y[\phi]$ is a version of $X(\phi)$. Now, as the topology ϑ_X is weaker than the nuclear topology on Φ , then Φ (equipped with the nuclear topology) is continuously embedded in Φ_{ϑ_X} , but as this last is continuously embedded in $\widetilde{\Phi_{\vartheta_X}}$, it follows that Φ is continuously embedded in $\widetilde{\Phi_{\vartheta_X}}$. Then, the space $(\widetilde{\Phi_{\vartheta_X}})'_\beta$ is continuously embedded in Φ'_β , and this implies that Y is a Φ'_β -valued, regular, continuous process that is a version of X .

Finally, to prove the uniqueness note that if Z is another version of X satisfying the properties in the Theorem, then for every $\phi \in \Phi$ and $t \geq 0$, $Y_t[\phi] = X_t(\phi) = Z_t(\phi)$ $\mathbb{P}$ -a.e., and because Y and Z are both continuous and regular Φ'_β -valued processes it follows from Proposition 2.12 that Y and Z are indistinguishable processes. $\square$

Remark 3.9. *It follows from the proof of Theorem 3.2 (in particular from the proof of Proposition 3.3) that the $(\widetilde{\Phi_{\vartheta_X}})'_\beta$ -valued continuous (respectively càdlàg) version Y of X can be chosen in such a way so that for every $\omega \in \Omega$ and $T > 0$ there exists a continuous Hilbertian semi-norm $\varrho = \varrho(\omega, T)$ on Φ such that the map $t \mapsto Y_t(\omega)$ is continuous (respectively càdlàg) from $[0, T]$ into the Hilbert space Φ'_ϱ .*

The assumption (2) in Theorem 3.2 of equicontinuity for every $T > 0$ of the family of cylindrical random variables $\{X_t : t \in [0, T]\}$ was important to show that the map $\phi \mapsto \{\widehat{X}_t(\phi)\}_{t \in [0, T]}$ from Φ into $C_T(\mathbb{R})$ (respectively $D_T(\mathbb{R})$) is continuous (Lemma 3.5). The next results shows that we can obtain the same conclusion without the above local equicontinuity property under the additional assumption that Φ is ultrabornological, i.e. that Φ is the inductive limit of Banach spaces (see [26], Theorem 13.2.11, p.448-9). Note that in the result below no nuclearity of the space is required.

Proposition 3.10. *Let Ψ be an ultrabornological space and let $X = \{X_t\}_{t \in [0, T]}$, be a cylindrical process in Ψ' such that for each $\psi \in \Psi$, the real-valued process $\{X_t(\psi)\}_{t \in [0, T]}$ has a continuous (respectively càdlàg) version. Assume that for every $t \in [0, T]$ the map $X_t : \Psi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is continuous. Then, the linear mapping from Ψ into $C_T(\mathbb{R})$ (respectively $D_T(\mathbb{R})$) given by $\psi \mapsto \{X_t(\psi)\}_{t \in [0, T]}$ is continuous.*

Proof. We can prove that the map $\psi \mapsto \{X_t(\psi)\}_{t \in [0, T]}$ from Ψ into $C_T(\mathbb{R})$ (respectively $D_T(\mathbb{R})$) is sequentially closed by following similar arguments to those used in the proof of Lemma 3.5. Then, as Ψ is ultrabornological and $C_T(\mathbb{R})$ (respectively $D_T(\mathbb{R})$) is a complete, metrizable, topological vector space, the closed graph theorem (see [15], Proposition 2, Section 5.2 and Theorem 2, Section 5.4, p.90,94) shows that $\psi \mapsto \{X_t(\psi)\}_{t \in [0, T]}$ is continuous. $\square$

We have the following version of the regularization theorem for cylindrical processes in the dual of an ultrabornological nuclear space.

Corollary 3.11. *Let Φ be an ultrabornological nuclear space. Let $X = \{X_t\}_{t \geq 0}$ be a cylindrical process in Φ' satisfying:*

- (1) *For each $\phi \in \Phi$, the real-valued process $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$ has a continuous (respectively càdlàg) version.*
- (2) *For every $t \geq 0$, $X_t : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is continuous.*

Then, there exists a countably Hilbertian topology ϑ_X on Φ and a $(\widetilde{\Phi_{\vartheta_X}})'_{\beta}$ -valued continuous (respectively càdlàg) process $Y = \{Y_t\}_{t \geq 0}$, such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \geq 0}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$. Moreover, Y is a Φ'_{β} -valued, regular, continuous (respectively càdlàg) version of X that is unique up to indistinguishable versions.

Proof. Let $T > 0$. With the same terminology as in the proof of Proposition 3.3, it follows from Proposition 3.10 that the linear mapping from Φ into $C_T(\mathbb{R})$ (respectively $D_T(\mathbb{R})$) given by $\psi \mapsto \{\hat{X}_t(\psi)\}_{t \in [0, T]}$ is continuous. We can use this result to prove Lemma 3.7. Then, the same arguments used in the remain of the proof of Proposition 3.3 show that the conclusions of Proposition 3.3 are still valid in our present context. Therefore, the proof of Theorem 3.2 can be replicated without any change and showing the existence of the countably Hilbertian topology θ_X and the $(\Phi_{\theta_X})'_{\beta}$ -valued process Y satisfying the conclusions of the corollary. $\square$

From the above result we obtain the following version of the regularization theorem for stochastic processes taking values in the dual of an ultrabornological nuclear space.

Corollary 3.12. *Let Φ be an ultrabornological nuclear space. Let $X = \{X_t\}_{t \geq 0}$ be a Φ'_{β} -valued process satisfying:*

- (1) *For each $\phi \in \Phi$, the real-valued process $X[\phi] = \{X_t[\phi]\}_{t \geq 0}$ has a continuous (respectively càdlàg) version.*
- (2) *For every $t \geq 0$, the distribution μ_{X_t} of X_t is a Radon probability measure.*

Then, there exists a countably Hilbertian topology ϑ_X on Φ and a $(\widetilde{\Phi_{\vartheta_X}})'_{\beta}$ -valued continuous (respectively càdlàg) process $Y = \{Y_t\}_{t \geq 0}$, such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \geq 0}$ is a version of $X[\phi] = \{X_t[\phi]\}_{t \geq 0}$. Moreover, Y is a Φ'_{β} -valued, regular, continuous (respectively càdlàg) version of X that is unique up to indistinguishable versions.

Proof. First, because the space Φ is ultrabornological it is also barreled (see [26], p.449). Hence, it follows from the assumption (2) in the Corollary and from Theorem 2.10 that for each $t \geq 0$ the map $X_t : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$, $\phi \mapsto X_t[\phi]$ is continuous. Then, the existence of the countably Hilbertian topology θ_X and the $(\Phi_{\theta_X})'_{\beta}$ -valued process Y satisfying the conclusions of the corollary follows from Corollary 3.11. $\square$

We finish this section with some comparisons of our results with those obtained by other authors.

First, it is clear that Theorem 3.2 is a generalization of the regularization theorem of Itô and Nawata (Theorem 2.9).

Now, if Φ is a nuclear space that is also a Fréchet space or the countable inductive limit of Fréchet spaces, then Φ is an ultrabornological space (see [15], Corollaries 4 and 5, Section 13.1, p.273). In this case if $X = \{X_t\}_{t \geq 0}$ is a Φ'_{β} -valued process then for every $t \geq 0$ we have that μ_{X_t} is a Radon probability measure (see Section 2.2). Then, if for each $\phi \in \Phi$, the real-valued process $X[\phi] = \{X_t[\phi]\}_{t \geq 0}$ has a continuous (respectively càdlàg) version, then Corollary 3.12 shows that X has a Φ'_{β} -valued continuous (respectively càdlàg) version. Therefore, Corollary 3.12 (and consequently Theorem 3.2) generalizes the results obtained by Mitoma [24], Fouque [10] and Fernique [8].

On the other hand, Martias [21] showed a version of the regularization theorem under the assumptions that Φ is a separable nuclear space and that X is of the form $X : [0, T] \times \Omega \rightarrow \Phi'$, such that for each $\phi \in \Phi$, $\{X_t[\phi]\}_{t \in [0, T]}$ is a real-valued process with a continuous (respectively càdlàg) version and also assuming that the map $\phi \mapsto \{X_t[\phi]\}_{t \in [0, T]}$ from Φ into $C_T(\mathbb{R})$ (respectively $D_T(\mathbb{R})$) is continuous. Note that the assumptions on Φ and X in Theorem 3.2 are weaker than the assumptions used by Martias. In particular, one can easily check that the continuity of the map

$\phi \mapsto \{X_t[\phi]\}_{t \in [0, T]}$ implies assumption (2) in Theorem 3.2. Hence, Theorem 3.2 constitutes a generalization of the result obtained by Martias.

4 Existence of Continuous and Càdlàg Versions in a Hilbert Space Continuously Embedded in Φ'_β

In this section we introduce conditions for the existence of continuous and càdlàg versions taking values in a Hilbert space Φ'_ϱ , where ϱ is a continuous Hilbertian semi-norm on Φ . We start with the following result that specializes the Regularization Theorem.

Theorem 4.1. *Let $X = \{X_t\}_{t \geq 0}$ be a cylindrical process in Φ' satisfying:*

- (1) *For each $\phi \in \Phi$, the real-valued process $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$ has a continuous (respectively càdlàg) version.*
- (2) *There exists a continuous Hilbertian semi-norm p on Φ such that for every $t \geq 0$, $X_t : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is p -continuous.*

Then, there exists a continuous Hilbertian semi-norm ϱ on Φ , $p \leq \varrho$, such that $i_{p, \varrho}$ is Hilbert-Schmidt and a Φ'_ϱ -valued continuous (respectively càdlàg) process $Y = \{Y_t\}_{t \geq 0}$, such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \geq 0}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$. Moreover, Y is unique up to indistinguishable versions in Φ'_β .

Proof. Let $T > 0$. With the same terminology as in the proof of Proposition 3.3, the assumption (2) of the Theorem implies that for every $t \in [0, T]$ the map $\hat{X}_t : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is p -continuous. Then, for all $t \in [0, T]$ the map $\hat{X}_t$ has a continuous and linear extension $\tilde{X}_t : \Phi_p \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$. Hence, from similar arguments to those in the proof of Lemma 3.5 we can show that the linear mapping $\tilde{X} : \Phi_p \rightarrow C_T(\mathbb{R})$, $\phi \mapsto \tilde{X}(\phi) = \{\tilde{X}_t(\phi)\}_{t \in [0, T]}$ is continuous. This in turns implies that the map $\hat{X} : \Phi \rightarrow C_T(\mathbb{R})$, $\phi \mapsto \hat{X}(\phi) = \{\hat{X}_t(\phi)\}_{t \in [0, T]}$ is p -continuous. Hence, in the proof of Lemma 3.8, we have that (3.3) is satisfied for $p_n = p$, for all $n \in \mathbb{N}$. In that case, we have in (3.12), (3.14) and (3.15) that $q_n = q$ for some continuous Hilbertian semi-norm q on Φ , $p \leq q$, and such that $i_{p, q}$ is Hilbert-Schmidt. Therefore, we have from (3.16) and (3.18) that $\Lambda_n = \Lambda_m$ and $Y^{(n)} = Y^{(m)}$ for all $m, n \in \mathbb{N}$. If we choose ϱ , $q \leq \varrho$, such that $i_{q, \varrho}$ is Hilbert-Schmidt, and if in the proof of Proposition 3.3 we take $\varrho_n = \varrho$ for all $n \in \mathbb{N}$, then $Y = \{Y_t\}_{t \in [0, T]}$ defined by (3.24) is a Φ'_ϱ -valued continuous processes such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \in [0, T]}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \in [0, T]}$.

Let $\{T_k\}_{k \in \mathbb{N}}$ an increasing sequence of positive numbers such that $\lim_{k \rightarrow \infty} T_k = \infty$. Then, as proved on the above paragraph for each $k \in \mathbb{N}$ there exists a Φ'_ϱ -valued continuous process $Y^{(k)} = \{Y_t^{(k)}\}_{t \in [0, T_k]}$, such that for every $\phi \in \Phi$, $\{Y_t^{(k)}[\phi]\}_{t \in [0, T_k]}$ is a version of $\{X_t(\phi)\}_{t \in [0, T_k]}$.

Take $Y = \{Y_t\}_{t \geq 0}$ defined by the prescription $Y_t = Y_t^{(k)}$ if $t \in [0, T_k]$. Note that Y is a well-defined Φ'_ϱ -valued process. To prove this, observe that for each $k \in \mathbb{N}$, $Y^{(k)}$ and $Y^{(k+1)}$ are continuous processes such that for every $\phi \in \Phi$, $Y_t^{(k)}[\phi] = X_t(\phi) = Y_t^{(k+1)}[\phi]$ $\mathbb{P}$ -a.e. for all $t \in [0, T_k]$, then Proposition 2.12 shows that $\{Y_t^{(k)}\}_{t \in [0, T_k]}$ and $\{Y_t^{(k+1)}\}_{t \in [0, T_k]}$ are indistinguishable processes. Therefore, Y is a Φ'_ϱ -valued continuous process such that for every $\phi \in \Phi$, $Y[\phi]$ is a version of $X(\phi)$. Moreover, from Proposition 2.12 again it follows that Y is unique up to indistinguishable versions in Φ'_β . $\square$

Now we consider conditions for the existence of continuous or càdlàg version such that in a given bounded interval of time they take values and have uniform finite moments in some Hilbert space Φ'_ϱ . For the proof we will need the following terminology. For $n \in \mathbb{N}$, we denote by $C_T^n(\mathbb{R})$ (respectively $D_T^n(\mathbb{R})$) the linear space

of all the continuous (respectively càdlàg) processes satisfying $\mathbb{E} \sup_{t \in [0, T]} |Z_t|^n < \infty$. It is a Banach space equipped with the norm $\|Z\|_{n, T} = \left(\mathbb{E} \sup_{t \in [0, T]} |Z_t|^n \right)^{1/n}$.

Theorem 4.2. *Let $X = \{X_t\}_{t \geq 0}$ be a cylindrical process in Φ' satisfying:*

- (1) *For each $\phi \in \Phi$, the real-valued process $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$ has a continuous (respectively càdlàg) version.*
- (2) *For every $T > 0$, the family $\{X_t : t \in [0, T]\}$ of linear maps from Φ into $L^0(\Omega, \mathcal{F}, \mathbb{P})$ is equicontinuous.*

- (3) *There exists $n \in \mathbb{N}$ such that $\forall T > 0$, $\mathbb{E} \left(\sup_{t \in [0, T]} |X_t[\phi]|^n \right) < \infty$, $\forall \phi \in \Phi$.*

Then, there exists a countably Hilbertian topology ϑ_X on Φ determined by an increasing sequence $\{q_k\}_{k \in \mathbb{N}}$ of continuous Hilbertian semi-norms on Φ and a $(\Phi_{\vartheta_X})'_\beta$ -valued continuous (respectively càdlàg) process $Y = \{Y_t\}_{t \geq 0}$, satisfying:

- (a) *For every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \geq 0}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$,*
- (b) *For every $T > 0$, there exists $k \in \mathbb{N}$ such that $\{Y_t\}_{t \in [0, T]}$ is a Φ'_{q_k} -valued continuous (respectively càdlàg) process satisfying $\mathbb{E} \left(\sup_{t \in [0, T]} q'_k(Y_t)^n \right) < \infty$.*

Furthermore, Y is a Φ'_β -valued, regular, continuous (respectively càdlàg) version of X that is unique up to indistinguishable versions.

Proof. We prove the continuous case as the càdlàg case follows from similar arguments. We do this in two steps.

Step 1 We are going to prove the following: *For every $T > 0$ there exists a continuous Hilbertian semi-norm q on Φ and a Φ'_q -valued continuous process $Y = \{Y_t\}_{t \in [0, T]}$, such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \in [0, T]}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \in [0, T]}$ and $\mathbb{E} \left(\sup_{t \in [0, T]} q'(Y_t)^n \right) < \infty$.*

Let $T > 0$ and define $p : \Phi \rightarrow [0, \infty)$ by

$$p(\phi) = \left[\mathbb{E} \left(\sup_{t \in [0, T]} |X_t(\phi)|^n \right) \right]^{1/n}, \quad \forall \phi \in \Phi. \quad (4.1)$$

We proceed to show that p is a continuous semi-norm on Φ . First, observe that the map from Φ into $C_T^n(\mathbb{R})$ given by $\phi \mapsto X(\phi) = \{X_t(\phi)\}_{t \in [0, T]}$ is linear. Then, because $\|\cdot\|_{n, T}$ is a norm on $C_T^n(\mathbb{R})$ and $p(\phi) = \|X(\phi)\|_{n, T}$ for all $\phi \in \Phi$, it follows that p is a semi-norm.

To prove that p is continuous, one can use the closed graph theorem and similar arguments to those used in the proof of Lemma 3.5 to show that the map $\phi \mapsto X(\phi)$ from Φ into $C_T^n(\mathbb{R})$ is continuous. But because $p(\phi) = \|X(\phi)\|_{n, T}$, for all $\phi \in \Phi$, then the continuity of the maps $\phi \mapsto X(\phi)$ and $Z \mapsto \|Z\|_{n, T}$ implies that p is continuous.

Now, from the Markov and Jensen inequalities, for any $\epsilon > 0$ we have

$$\mathbb{P} \left(\omega : \sup_{t \in [0, T]} |X_t(\phi)(\omega)| > \epsilon \right) \leq \frac{1}{\epsilon} \left[\mathbb{E} \left(\sup_{t \in [0, T]} |X_t(\phi)|^n \right) \right]^{1/n} = \frac{1}{\epsilon} p(\phi), \quad \forall \phi \in \Phi.$$

Therefore, the map $X : \Phi \rightarrow C_T(\mathbb{R})$ given by $\psi \mapsto X(\psi) = \{X_t(\psi)\}_{t \in [0, T]}$ is p -continuous. Now, as the topology on Φ is generated by a family of Hilbertian semi-norms and because the semi-norm p is continuous, there exists a continuous Hilbertian semi-norm ϱ on Φ such that $p \leq \varrho$. This implies that the map $X : \Phi \rightarrow C_T(\mathbb{R})$ is ϱ -continuous. Then, it follows from Theorem 4.1 that there exists a continuous Hilbertian semi-norm r on Φ , $\varrho \leq r$ such that $i_{r, \varrho}$ is Hilbert-Schmidt and such that X has a Φ'_r -valued continuous version $\hat{Y} = \{\hat{Y}_t\}_{t \in [0, T]}$.

Let q be a continuous Hilbertian semi-norm on Φ such that $r \leq q$ and $i_{r, q}$ is Hilbert-Schmidt. Then, the dual operator $i'_{r, q} : \Phi'_r \rightarrow \Phi'_q$ is Hilbert-Schmidt and hence is n -summing (see [7], Corollary 4.13, p.85). Then, from the Pietsch

domination theorem (see [7], Theorem 2.12, p.44) there exists a constant $C > 0$, and a Radon probability measure ν on the unit ball $B_r^*(1)$ of Φ_r (equipped with the weak topology) such that,

$$q'(i'_{r,q}f) \leq C \cdot \left(\int_{B_r^*(1)} |f[\phi]|^n \nu(d\phi) \right)^{1/n}, \quad \forall f \in \Phi'_r. \quad (4.2)$$

As $\widehat{Y}$ is a Φ'_r -valued continuous process, then $\phi \mapsto \widehat{Y}[\phi] = \{\widehat{Y}_t[\phi]\}_{t \in [0,T]}$ is a continuous and linear map from Φ_r into $C_T^n(\mathbb{R})$. Therefore, it follows from (4.2) that,

$$\begin{aligned} \mathbb{E} \left(\sup_{t \in [0,T]} q'(i'_{r,q} \widehat{Y}_t)^n \right) &\leq C^n \mathbb{E} \sup_{t \in [0,T]} \int_{B_r^*(1)} |\widehat{Y}_t[\phi]|^n \nu(d\phi) \\ &\leq C^n \int_{B_r^*(1)} \left\| \widehat{Y}[\phi] \right\|_{c,n,T}^n \nu(d\phi) \\ &\leq C^n \left\| \widehat{Y} \right\|_{\mathcal{L}(\Phi_r, C_T^n(\mathbb{R}))}^n < \infty. \end{aligned}$$

Hence, $Y = \{Y_t\}_{t \in [0,T]}$, defined by $Y_t = i'_{r,q} \widehat{Y}_t$ for every $t \in [0,T]$, is a Φ'_q -valued continuous process such that for every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \in [0,T]}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \in [0,T]}$ and such that $\mathbb{E} \left(\sup_{t \in [0,T]} q'(Y_t)^n \right) < \infty$.

Step 2 Let $\{T_k\}_{k \in \mathbb{N}}$ an increasing sequence of positive numbers such that $\lim_{k \rightarrow \infty} T_k = \infty$. Then, it follows from Step 1 that for each $k \in \mathbb{N}$ there exists a continuous Hilbertian semi-norm q_k on Φ and a Φ'_{q_k} -valued continuous process $Y^{(k)} = \{Y_t^{(k)}\}_{t \in [0,T_k]}$, such that for every $\phi \in \Phi$, $Y^{(k)}[\phi] = \{Y_t^{(k)}[\phi]\}_{t \in [0,T_k]}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \in [0,T_k]}$ and $\mathbb{E} \left(\sup_{t \in [0,T_k]} q'_k(Y_t^{(k)})^n \right) < \infty$.

Without loss of generality we can assume that the sequence of semi-norms $\{q_k\}_{k \in \mathbb{N}}$ is increasing. Let ϑ_X be the countably Hilbertian topology on Φ generated by the semi-norms $\{q_k\}_{k \in \mathbb{N}}$. Take $Y = \{Y_t\}_{t \geq 0}$ defined by the prescription $Y_t = Y_t^{(k)}$ if $t \in [0, T_k]$. In a similar way as we did in the proof of Theorem 3.2, from the corresponding properties of the processes $Y^{(k)}$ s we can conclude that Y is a $(\Phi_{\vartheta_X})'_\beta$ -valued continuous process satisfying the properties (a) and (b) of the Theorem. Finally, as in the proof of Theorem 3.2 we can show that as a Φ'_β -valued process Y is unique up to indistinguishable versions. $\square$

The following result is an specialized version of Theorem 4.2 that establish conditions for the existence of continuous or càdlàg version in single Hilbert space Φ'_ϱ with uniform finite moments in every bounded interval of time.

Theorem 4.3. *Let $X = \{X_t\}_{t \geq 0}$ be a cylindrical process in Φ' satisfying:*

- (1) *For each $\phi \in \Phi$, the real-valued process $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$ has a continuous (respectively càdlàg) version.*
- (2) *There exists $n \in \mathbb{N}$ and a continuous Hilbertian semi-norm ϱ on Φ such that for all $T > 0$ there exists $C(T) > 0$ such that*

$$\mathbb{E} \left(\sup_{t \in [0,T]} |X_t(\phi)|^n \right) \leq C(T) \varrho(\phi)^n, \quad \forall \phi \in \Phi. \quad (4.3)$$

Then, there exists a continuous Hilbertian semi-norm q on Φ , $\varrho \leq q$, such that $i_{\varrho,q}$ is Hilbert-Schmidt and there exists a Φ'_q -valued continuous (respectively càdlàg) process, satisfying:

- (a) *For every $\phi \in \Phi$, $Y[\phi] = \{Y_t[\phi]\}_{t \geq 0}$ is a version of $X(\phi) = \{X_t(\phi)\}_{t \geq 0}$,*
- (b) *For every $T > 0$, $\mathbb{E} \left(\sup_{t \in [0,T]} q'(Y_t)^n \right) < \infty$.*

Furthermore, Y is a Φ'_β -valued continuous (respectively càdlàg) version of X that is unique up to indistinguishable versions.

Proof. As before, we will prove the continuous case as the càdlàg case follows similarly.

Fix $T > 0$. Note that (4.3) is equivalent to $p \leq C(T)^{1/n} \varrho$, where p is the semi-norm defined in (4.1). Therefore, the map from Φ into $C_T(\mathbb{R})$ given by $\phi \mapsto X(\phi) = \{X_t(\phi)\}_{t \in [0, T]}$ is ϱ -continuous. Hence, similar arguments to those in Step 1 of the proof of Theorem 4.2 show that there exists a continuous Hilbertian semi-norm q on Φ (only depending on ϱ), $\varrho \leq q$, such that $i_{\varrho, q}$ is Hilbert-Schmidt, and a Φ'_q -valued continuous version $Y = \{Y_t\}_{t \in [0, T]}$ of $\{X_t\}_{t \in [0, T]}$ satisfying $\mathbb{E} \left(\sup_{t \in [0, T]} q'(Y_t)^n \right) < \infty$.

Let $\{T_k\}_{k \in \mathbb{N}}$ an increasing sequence of positive numbers such that $\lim_{k \rightarrow \infty} T_k = \infty$. Then, as proved in the above paragraph for each $k \in \mathbb{N}$ there exists a Φ'_q -valued continuous version $Y^{(k)} = \{Y_t^{(k)}\}_{t \in [0, T_k]}$ of $\{X_t\}_{t \in [0, T_k]}$ that satisfies $\mathbb{E} \left(\sup_{t \in [0, T_k]} q'(Y_t^{(k)})^n \right) < \infty$. Observe that for each $k \in \mathbb{N}$, the Φ'_q -valued continuous processes $\{Y_t^{(k)}\}_{t \in [0, T_k]}$ and $\{Y_t^{(k+1)}\}_{t \in [0, T_k]}$ are indistinguishable. Therefore, if we define $Y = \{Y_t\}_{t \geq 0}$ by the prescription $Y_t = Y_t^{(k)}$ if $t \in [0, T_k]$, then Y is a Φ'_q -valued continuous version of X satisfying $\mathbb{E} \left(\sup_{t \in [0, T]} q'(Y_t)^n \right) < \infty$, for all $T > 0$. $\square$

5 Applications to Cylindrical Martingales in Φ'

In this section we apply the results in Section 4 to study some properties of cylindrical martingales in Φ' . Consider a filtration $\{\mathcal{F}_t\}_{t \geq 0}$ on $(\Omega, \mathcal{F}, \mathbb{P})$ that satisfies the *usual conditions*, i.e. it is right continuous and $\mathcal{F}_0$ contains all sets of $\mathcal{F}$ of $\mathbb{P}$ -measure zero.

Let $n \geq 1$. Denote by $\mathcal{M}^n(\mathbb{R})$ the space of all the real-valued martingales $M = \{M_t\}_{t \geq 0}$ satisfying

$$\|M\|_{\mathcal{M}^n(\mathbb{R})} = \left[\mathbb{E} \left(\sup_{t \geq 0} |M_t|^n \right) \right]^{1/n} < \infty. \quad (5.1)$$

It is clear that $\|\cdot\|_{\mathcal{M}^n(\mathbb{R})}$ defines a norm on $\mathcal{M}^n(\mathbb{R})$. Moreover, $(\mathcal{M}^n(\mathbb{R}), \|\cdot\|_{\mathcal{M}^n(\mathbb{R})})$ is a Banach space (see [6]). Note that if the real-valued martingale $M = \{M_t\}_{t \geq 0}$ satisfies $\sup_{t \geq 0} \mathbb{E}(|M_t|^n) < \infty$, for $n > 2$, then Doob's inequality shows that $\|M\|_{\mathcal{M}^n(\mathbb{R})} < \infty$ and hence $M \in \mathcal{M}^n(\mathbb{R})$.

Definition 5.1. A *cylindrical martingale* in Φ' is a cylindrical process $M = \{M_t\}_{t \geq 0}$ such that for each $\phi \in \Phi$, the real-valued process $M(\phi) = \{M_t(\phi)\}_{t \geq 0}$ is a $\{\mathcal{F}_t\}$ -adapted martingale.

The following theorem contains some of the basic properties of cylindrical martingales in Φ' .

Theorem 5.2. Let $M = \{M_t\}_{t \geq 0}$ be a cylindrical martingale in Φ' such that for each $t \geq 0$ the map $M_t : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is continuous. Then, M has a Φ'_β -valued càdlàg version $\widetilde{M} = \{\widetilde{M}_t\}_{t \geq 0}$. Moreover, we have the following:

- (1) If M is n -th integrable, for $n \geq 2$, then for each $T > 0$ there exists a continuous Hilbertian semi-norm q_T on Φ such that $\{\widetilde{M}_t\}_{t \in [0, T]}$ is a Φ'_{q_T} -valued càdlàg martingale satisfying $\mathbb{E} \left(\sup_{t \in [0, T]} q'_T(\widetilde{M}_t)^n \right) < \infty$.
- (2) Moreover, if for $n \geq 2$, $\sup_{t \geq 0} \mathbb{E}(|M_t(\phi)|^n) < \infty$ for each $\phi \in \Phi$, then there exists a continuous Hilbertian semi-norm q on Φ such that $\widetilde{M}$ is a Φ'_q -valued càdlàg martingale satisfying $\mathbb{E} \left(\sup_{t \geq 0} q'(\widetilde{M}_t)^n \right) < \infty$.

If for each $\phi \in \Phi$ the real-valued process $\{M_t(\phi)\}_{t \geq 0}$ has a continuous version, then $\widetilde{M}$ can be chosen to be continuous and such that it satisfies (1)–(2) above replacing the property càdlàg by continuous.

Proof. To show the existence of the Φ'_β -valued càdlàg version $\widetilde{M}$ of M we need to check that the conditions in Theorem 3.2 are satisfied. As for each $\phi \in \Phi$, $M_t(\phi)$ is a real-valued martingale, then it has a càdlàg version. Then it only remains to verify that for every $T > 0$ the family $\{M_t : t \in [0, T]\} \subseteq \mathcal{L}(\Phi, L^0(\Omega, \mathcal{F}, \mathbb{P}))$ is equicontinuous.

Let $T > 0$. First note that because for all $\phi \in \Phi$, $\{|M_t(\phi)|\}_{t \geq 0}$ is a positive submartingale, then by Doob's tail martingale inequality we have for any $\epsilon > 0$,

$$\mathbb{P} \left(\sup_{t \in [0, T]} |M_t(\phi)| > \epsilon \right) \leq \frac{1}{\epsilon} \mathbb{E}(|M_T(\phi)|). \quad (5.2)$$

Define $p_T : \Phi \rightarrow [0, \infty)$ by $p_T(\phi) = \mathbb{E}(|M_T(\phi)|)$, for all $\phi \in \Phi$. Then, p_T is a semi-norm on Φ . Moreover, from the continuity of the map $M_T : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ and by following similar ideas to those used in the proof of Lemma 3.4, we can show that there exists a countably Hilbertian topology θ on Φ such that the map M_T is θ -continuous. By means of an application of the closed graph theorem as in the proof of Lemma 3.5 we can show that M_T is continuous as a map from Φ_θ into $L^1(\Omega, \mathcal{F}, \mathbb{P})$. This in turns implies that p_T is continuous. Then, it follows from (5.2) that the family $\{M_t : t \in [0, T]\} \subseteq \mathcal{L}(\Phi, L^0(\Omega, \mathcal{F}, \mathbb{P}))$ is equicontinuous. Therefore, the existence of the Φ'_β -valued càdlàg version $\widetilde{M}$ of M is a consequence of Theorem 3.2.

If M is n -th integrable, for $n \geq 2$, then it follows from Doob's inequality that for every $T > 0$ and $\phi \in \Phi$ we have

$$\mathbb{E} \left(\sup_{t \in [0, T]} |M_t(\phi)|^n \right) \leq m^n \mathbb{E}(|M_T(\phi)|^n) < \infty,$$

where $1/n + 1/m = 1$. Then, (1) follows from Theorem 4.2.

To prove (2), note that for each $\phi \in \Phi$ the condition $\sup_{t \geq 0} \mathbb{E}(|M_t(\phi)|^n) < \infty$ implies that $M(\phi) = \{M_t(\phi)\}_{t \geq 0} \in \mathcal{M}^n(\mathbb{R})$. Then, M defines a linear map $\phi \mapsto M(\phi)$ from Φ into $\mathcal{M}^n(\mathbb{R})$. We can show that this map is continuous by using similar arguments to those used in Lemma 3.5. Then, it follows from (5.1) that $p : \Phi \rightarrow [0, \infty)$ defined by

$$p(\phi) = \left[\mathbb{E} \left(\sup_{t \geq 0} |M_t[\phi]|^n \right) \right]^{1/n}, \quad \forall \phi \in \Phi,$$

is a continuous semi-norm on Φ . Let ϱ be a continuous Hilbertian semi-norm on Φ such that $p \leq \varrho$. Then, it follows from the definition of p and Chebyshev's inequality that for every $\epsilon > 0$ we have

$$\mathbb{P} \left(\sup_{t \geq 0} |M_t(\phi)| > \epsilon \right) \leq \frac{1}{\epsilon^n} p(\phi)^n \leq \frac{1}{\epsilon^n} \varrho(\phi)^n, \quad \forall \phi \in \Phi.$$

Therefore, for every $t \geq 0$ the map $M_t : \Phi \rightarrow L^0(\Omega, \mathcal{F}, \mathbb{P})$ is ϱ -continuous. Hence, Theorem 4.1 shows that there exists a continuous Hilbertian semi-norm r on Φ , $\varrho \leq r$ such that $i_{r, \varrho}$ is Hilbert-Schmidt and such that M has a Φ'_r -valued càdlàg version $\widetilde{M} = \{\widetilde{M}_t\}_{t \geq 0}$ that is a martingale. Moreover, because $p \leq r$ is clear that $\widetilde{M}$ defines a continuous and linear map $\phi \mapsto \widetilde{M}(\phi)$ from Φ_r into $\mathcal{M}^n(\mathbb{R})$. Then, from a modification of the arguments in Step 2 of the proof of Theorem 4.2, we can show that there exists a continuous Hilbertian semi-norm q on Φ , $r \leq q$, such that M has a Φ'_q -valued càdlàg version $\widetilde{M} = \{\widetilde{M}_t\}_{t \geq 0}$ that is a martingale and satisfies $\mathbb{E} \left(\sup_{t \geq 0} q'(\widetilde{M}_t)^n \right) < \infty$. $\square$

Remark 5.3. *The results in Theorem 5.2 (for the case $n = 2$) were originally proved by Mitoma (see [23]; see also [17], Theorems 3.1.3-4) for the case of martingales in the strong dual of a nuclear Fréchet space. Note that we have been able to extend these results to any nuclear space and for n -integrable martingales for any $n \geq 2$.*

Acknowledgements I want to express my deepest gratitude to my PhD Supervisor David B. Applebaum for all his advice, comments and suggestions that contributed to improve this work. Also, I would like to thank both the School of Mathematics and Statistics (SoMaS) of The University of Sheffield and the Office of International Affairs and External Cooperation (OAICE) of The University of Costa Rica for providing financial support.

References

- [1] D. Applebaum and M. Riedle, Cylindrical Lévy processes in Banach spaces, *Proc. London Math. Soc.*, 101(3): 697–726, 2010.
- [2] A. Badrikian, *Séminaire Sur les Fonctions Aléatoires Linéaires et les Mesures Cylindriques*, Lecture Notes in Math. 139, Springer, 1970.
- [3] T. Bojdecki and J. Jakubowski, Stochastic Integration for Inhomogeneous Wiener Process in the Dual of a Nuclear Space, *J. Multivariate Anal.*, 34:185–210, 1990.
- [4] T. Bojdecki and A. Talarczyk, Particle picture approach of self-intersection local time for density processes in $\mathcal{S}'(\mathbb{R}^d)$, *Stoch. Proc. Appl.*, 115:449–479, 2005.
- [5] Yu. L. Dalecky and S. V. Fomin, *Measure and Differential Equations in Infinite-Dimensional Space*, Mathematics and Its Applications 76, Springer Science+Business Media, 1991.
- [6] C. Dellacherie and P. A. Meyer, *Probabilities and Potential B. Theory of Martingales*, North-Holland Publishing Company, 1982.
- [7] J. Diestel and H. Jarchow and A. Tonge, *Absolutely Summing Operators*, Cambridge Series in Advanced Mathematics 43, Cambridge University Press, 1995.
- [8] X. Fernique, Fonctions aléatoires à valeurs dans certains espaces nucléaires, *Probab. Th. Rel. Fields*, 83:87–99, 1989.
- [9] C. A. Fonseca-Mora, *Stochastic Analysis with Lévy Noise in the Dual of a Nuclear Space*, PhD Thesis, The University of Sheffield, 2015.
- [10] J-P. Fouque, La convergence en loi pour les processus à valeurs dans un espace nucléaire, *Ann. Inst. H. Poincaré*, 20(3):225–245, 1984.
- [11] I. Gel’fand and N. Vilenkin, *Generalized Functions. Volume. 4. Applications to Harmonic Analysis*, Academic Press, 1964.
- [12] H. Heyer, *Structural Aspects in the Theory of Probability*, Series on Multivariate Analysis, World Scientific, second edition, 2010.
- [13] K. Itô, *Foundations of Stochastic Equations in Infinite Dimensional Spaces*, Series in Mathematics, SIAM, 1984.
- [14] K. Itô and M. Nawata, Regularization of linear random functionals, In *Probability Theory and Mathematical Statistics. Proceedings of the Fourth USSR-Japan Symposium, Tbilisi, USSR, August 23-29, 1982*, Lecture Notes in Math. 1021, Springer, 1983, 257–267.
- [15] H. Jarchow, *Locally Convex Spaces*, Mathematische Leitfäden, Springer, 1981.
- [16] G. Kallianpur and J. Xiong, Diffusion approximation of nuclear space-valued stochastic differential equations driven by Poisson random measures, *Ann. Appl. Probab.*, 5(2):493–517, 1995.

- [17] G. Kallianpur and J. Xiong, *Stochastic Differential Equations in Infinite Dimensional Spaces*, Lecture Notes-Monograph Series, Institute of Mathematical Statistics, 1995.
- [18] H. Kaspi and K. Ramanan, SPDE limits of many-server queues, *Ann. Appl. Probab.*, 23(1):145–229, 2013.
- [19] M. Ledoux and M. Talagrand, *Probability in Banach Spaces. Isoperimetry and Processes*, Ergebnisse der Mathematik und ihrer Grenzgebiete, Springer, 1991.
- [20] W. Linde, *Probability in Banach Spaces. Stable and Infinitely Divisible Distributions*, Wiley, 1986.
- [21] C. Martias, Sur les supports des processus à valeurs dans des espaces nucléaires, *Ann. Inst. H. Poincaré*, 24(3): 345–365, 1988.
- [22] R. A. Minlos, Generalized random processes and their extension to a measure, In *Selected Translations in Mathematical Statistics and Probability, Vol.3*, American Mathematical Society, 1963, 291-313.
- [23] I. Mitoma, Martingales of random distributions, *Mem. Fac. Sci. Kyushu Univ. Ser. A*, 35(1):185–197, 1981.
- [24] I. Mitoma, On the sample continuity of $\mathcal{S}'$ -processes, *J. Math. Soc. Japan*, 35(4):629–636, 1983.
- [25] I. Mitoma and S. Okada and Y. Okazaki, Cylindrical σ -algebra and cylindrical measure, *Osaka J. Math.*, 14(3):635-647, 1977.
- [26] L. Narici and E. Beckenstein, *Topological Vector Spaces*, Pure and Applied Mathematics, CRC Press, second edition, 2011.
- [27] S. Peszat and J. Zabczyk, *Stochastic Partial Differential Equations with Lévy Noise*, Encyclopedia of Mathematics and Its Applications, Cambridge University Press, 2007.
- [28] A. Pietsch, *Nuclear Locally Convex Spaces*, Ergebnisse der Mathematikund ihrer Grenzgebiete, Springer, 1972.
- [29] E. Priola and J. Zabczyk, Structural properties of semilinear SPDEs driven by cylindrical stable processes, *Probab. Theory Relat. Fields*, 149(1-2):97–137, 2011.
- [30] M. Riedle, Stochastic integration with respect to cylindrical Lévy processes in Hilbert spaces: an L^2 approach, *Infin. Dimens. Anal. Quantum. Probab. Relat. Top.*, 17: 2014.
- [31] M. Riedle, Ornstein-Uhlenbeck processes driven by cylindrical Lévy processes, *Potential Anal.*, 42:809–838, 2015.
- [32] V. V. Sazonov, *A Remark on Characteristic Functionals*, *Theory Probab. Appl.*, 3:188-192, 1958.
- [33] H. Schaefer, *Topological Vector Spaces*, Graduate Texts in Mathematics, Springer, second edition, 1999.
- [34] L. Schwartz, *Radon Measures on Arbitrary Topological Spaces and Cylindrical Measures*, Tata Institute of Fundamental Research Studies in Mathematics, Oxford University Press, 1973.
- [35] F. Trèves, *Topological Vector Spaces, Distributions and Kernels*, Pure and Applied Mathematics, Academic Press, 1967.
- [36] N. N. Vakhania and V. I. Tarieladze and S. A. Chobanyan, *Probability Distributions on Banach Spaces*, Reidel Publishing, 1987.
- [37] J. M. A. M. van Neerven and M.C. Veraar and L. Weis, Stochastic evolution equations in UMD Banach spaces, *Stoch. Process. Appl.*, 255(4):940–993, 2008.